

Cystic Meningioma with Interesting Imaging Characteristics in Frontal Region: A Case Report

Leila Aghaghazvini^{1*}, Milad Sanginabadi², Bahman Rasuli³, Salma Sefidbakht⁴ and Ala Torabi²

¹Associate Professor of Radiology, Department of Radiology, Shariati Hospital, Tehran University of Medical Science, Tehran, Iran

²Resident of Radiology, Department of Radiology, Shariati Hospital, Tehran University of Medical Science, Tehran, Iran

³Radiologist, Department of Radiology, Tehran University of Medical Science, Tehran, Iran

⁴Assistant Professor of Pathology, Department of Pathology, School of Medicine, Shariati Hospital, Tehran University of Medical Science, Tehran, Iran

***Corresponding Author:** Leila Aghaghazvini, Associate Professor of Radiology, Department of Radiology, Tehran University of Medical Science, Tehran, Iran.

Received: March 24, 2021; **Published:** March 29, 2021

Abstract

Introduction: Cystic meningioma is an unusual shape of meningioma and the imaging manifestations and site of the solid/cystic components of the mass increasing our difficulties to make a diagnosis.

Case Report: We report a case of a 78-year-old right-handed woman who presented with right-sided hemiparesis, urinary incontinuity and Broca aphasia. A large left-sided frontal region cystic extra-axial mass with a suspicious small solid nodule and adjacent bony erosion were detected on non-contrast CT of the brain. Brain MRI with gadolinium depicted the cystic and solid components exactly. A probable diagnosis of cystic meningioma was made and subsequently confirmed by histological study after surgical resection of the tumor. To confirm the category and subtypes of the tumor, histopathological examination of the tumor cells should be done.

Conclusion: Cystic or solid cystic intracranial lesions have several differential diagnoses making the final diagnosis difficult, in some cases. There are four types of cystic meningiomas according to Nauta classification that was revised by Jung, *et al.* Determination of the type of cystic meningioma is crucial for surgical planning. To confirm the category and subtypes of the tumor, histopathological examination of the tumor cells should be done. Our case was a type IV cystic meningioma with a small mural enhancing solid nodule but associated adjacent bony invasion.

Keywords: Cystic Meningioma; Imaging; MRI; Radiologic; Histopathologic

Background

Meningiomas are the most common primary brain tumors accounting for 15 - 30% of all intracranial tumors [1]. Meningioma with the cystic component is an uncommon type of meningiomas accounting for 4 - 7% of all intracranial meningiomas and frequently affects men and more commonly is found at frontoparietal areas; they could be misdiagnosed as a glioma or metastasis due to their radiologic appearance. Moreover, previous strokes, subdural hematomas, and abscess formation could be the other less common differential diagnosis [1-5].

Here we report a rare and interesting case of cystic meningioma that would easily cause misguidance in management plans and post-operative diagnosis.

Case Presentation

A 78-year-old right-handed woman who was evaluated for right-sided hemiparesis and urinary incontinency from three months and Broca aphasia since one month ago.

The initial lab test was in the normal range.

A well-defined more dominant extra-axial cystic mass in the left frontal lobe with peripheral mural nodule and adjacent bone changes was observed in the initial unenhanced CT scan (Figure 1). On MRI an extra-axial cystic mass with a peripheral mural nodule that showed avid enhancement follow contrast injection was detected. No surrounding edema or dural tail was found (Figure 2).




Figure 1: Brain CT without contrast.

*A: Brain windows, B: Bone windows. There is a 55*40 mm extra-axial hypodense cystic mass with a peripherally located isodense mural nodule in the left frontal region associated with adjacent frontal bone pressure erosion and minimal sclerosis. Mild mass effect and contra-lateral midline shift also are evident. No obvious edema is seen.*




Figure 2: A: T1 weighted image (axial), B: T2 weighted image (coronal), C: Fluid-attenuated inversion recovery (axial), D and E: post-contrast T1 weighted image (axial).

*There is a 40 * 55 * 60 mm cystic extra-axial mass with homogeneously enhancing mural nodule (15*15 mm) in the left frontal region with adjacent bone extension and irregularity and small enhancing component. Mass effect, contra-lateral midline shift, and subfalcial hernia are also depicted. No obvious edema or dural tail is seen.*

Our first diagnosis was a cystic meningioma, although the metastasis and hemangiopericytoma were included in our differential diagnosis regarding the extra-axial location of the mass with adjacent bony erosion.

The mass was completely resected and cystic meningioma was proved after histopathological study. The patient was discharged from our hospital with a good clinical condition 10 days after surgery. In the six months later follow-up, no recurrence was seen.




Figure 3: Microscopic evaluation of mass demonstrates a cellular neoplastic tissue composed of round to oval cells with indistinct cell membranes, eosinophilic cytoplasm and round uniform nuclei with fine chromatin and inconspicuous nucleoli. Some of them forming whorls. No mitotic figure or necrosis is noted.




Figure 4: Microscopic examination of the brain tumor cyst wall show strips of fibro-connective tissue lined by a single layer as well as proliferated meningotheelial cells that have PR and EMA positivity.

Discussion

About 2 - 3% of the population has an incidental asymptomatic meningioma and its incidence increases progressively with age [6]. Multiple risk factors including history of radiation (therapeutic or incidentally), head trauma, and genetic (Neurofibromatosis type 2) are suggested to contribute to the development of meningioma while most meningiomas are idiopathic with unknown etiology.

Meningiomas are extra-axial, often benign lesions that usually arise from meningotheelial cells of dural coverings of the brain and most of them arise from dural reflections and uncommonly may be found at optic nerve sheath and choroid plexus (intraventricular meningioma) [7]. Actually, meningioma is a typically solid tumor, and cystic meningioma is a rare form with an incidence of about 1.6 - 11.7% of meningiomas according to the previous studies [8]. Cystic meningiomas are usually located at cerebral convexity, mostly in the fronto-parietal region [2].

The exact mechanism of cyst formation in meningiomas is unknown, however, cellular necrosis, ischemia, hemorrhage as well as an active fluid secretion by the functional tumor cells into the mass are proposed as the underlying mechanism [9-11].

Intracranial lesions with cystic or solid cystic appearance as high-grade gliomas, metastasis, hemangiopericytoma, hydatid cyst and arachnoid cysts are important differential diagnoses leading to incorrect preoperative diagnosis. The differentiation may be difficult and challenging in some cases in which other modalities like brain angiography, DWI, and ADC images and MR spectrography could be helpful [3,9-12].

Classification of cystic meningioma was initially proposed by Nauta, *et al.* into four subtypes according to the location of the cyst to the brain and meningioma as followed:

- Type I: Intratumoral cysts that are centrally located in the tumor.
- Type II: Intratumoral cysts that are peripherally located and surrounded by meningioma.
- Type III: Peritumoral cyst in the adjacent parenchyma.
- Type IV: Peritumoral cyst between the tumor and the adjacent parenchyma [10].

The above classification was revised by Jung *et al* by adding type V as a combination of types I and III meningiomas [2].

The classification of cystic meningiomas is crucial for preoperative planning, especially for type II and III cysts because resection of the whole capsule with the cyst is necessary for type II while tumor excision only without the cyst is the choice plan in type III.

Nevertheless, discrimination of these two types is not always easy through MRI and so histological evaluation or microsurgical assessment of the cyst wall is essential [8,13].

The recent patient's MRI had depicted the presence of a Nauta type IV cystic meningioma. Widening of subarachnoid space, cerebrospinal fluid space mechanical trapping, and compression by the tumor and less frequently a true arachnoid cyst are the mechanisms of formation of peritumoral cyst of Nauta type IV [10]. Although this case was a type 4 cystic meningioma, it presented with a large cystic component and an enhancing small mural nodule, that leading to adjacent bone erosion and sclerosis and focally bony invasion of soft tissue as small enhancing component that most of the previous reported ones had a larger peripheral located soft tissue component leading to adjacent bony changes.

It should be mentioned that approximately 8% and 12% of all meningiomas with cystic changes are malignant and angioblastic respectively [10].

Management of cystic meningiomas like solid types is depending on signs, symptoms, and comorbidities of patients as well as the size and location of the tumor.

Small asymptomatic cystic meningiomas can be managed conservatively with serial imaging similar to the solid types. The recurrence rate after complete resection with or without embolization is about 6% as well [14,15].

Conclusion

Cystic meningiomas are uncommon forms that usually makes a dilemma in diagnosis and MRI diagnostic role is valuable for the diagnosis and classification of the lesions and finally, histopathological diagnosis is always required to confirm the diagnosis.

Funding

The authors have no funding to report.

Conflict of Interest

The authors have declared they have no competing interests.

Acknowledgement

Very thanks to the colleagues in Shariati hospital and the patient for cooperation in data collection.

Bibliography

1. Sotiriadis C., *et al.* "Cystic meningioma: diagnostic difficulties and utility of MRI in diagnosis and management". *BMJ Case Reports* (2015).
2. Fortuna A., *et al.* "Cystic meningiomas". *Acta Neurochirurgica* 90.1 (1988): 23-30.
3. Go KO., *et al.* "Cystic Meningiomas: Correlation between Radiologic and Histopathologic Features". *Brain Tumor Research and Treatment* 6.1 (2018): 13-21.
4. Guan TK., *et al.* "Infratentorial benign cystic meningioma mimicking a hemangioblastoma radiologically and a pilocytic astrocytoma intraoperatively: a case report". *Journal of Medical Case Reports* 7.1 (2013): 87.
5. Worthington C., *et al.* "Meningioma cysts". *Neurology* 35.12 (1985): 1720.
6. Nakasu S., *et al.* "Incidental meningiomas in autopsy study". *Surgical Neurology* 27.4 (1987): 319-322.
7. Watts J., *et al.* "Magnetic resonance imaging of meningiomas: a pictorial review". *Insights Imaging* 5.1 (2014): 113-122.
8. Go KO., *et al.* "Cystic Meningiomas: Correlation between Radiologic and Histopathologic Features". *Brain Tumor Research and Treatment* 6.1 (2018): 13-21.

9. Buetow MP, *et al.* "Typical, atypical, and misleading features in meningioma". *Radiographics* 11.6 (1991): 1087-1106.
10. Maj B, *et al.* "[Cystic meningioma: report of three cases]". *Neurologia i Neurochirurgia Polska* 36.1 (2002): 199-206.
11. Saxena D, *et al.* "MRI findings of an atypical cystic meningioma-A rare".
12. Jorge D, *et al.* "Cystic meningioma simulating arachnoid cyst: report of an unusual case". *Case Reports in Radiology* (2014).
13. Nauta HJ, *et al.* "Xanthochromic cysts associated with meningioma". *Journal of Neurology, Neurosurgery, and Psychiatry* 42.6 (1979): 529-535.
14. Ferrante L, *et al.* "MRI in the diagnosis of cystic meningiomas: Surgical implications". *Acta Neurochirurgica* 139.1 (1997): 8-11.
15. Mittal A, *et al.* "Cystic meningioma: unusual imaging appearance of a common intracranial tumor". *Proc* 23.4 (2010): 429-431.

Volume 4 Issue 4 April 2021

©All rights reserved by Leila Aghaghazvini, *et al.*