

Gastroesophageal Reflux Disease with Only Extraesophageal Symptoms. Case Report

Michael D Levin*

Department of Pediatric Radiology of the 1-st State Hospital, Minsk, Belarus and Dorot. Medical Center for Rehabilitation and Geriatrics, Netanya, Israel

***Corresponding Author:** Michael D Levin, Department of Pediatric Radiology of the 1-st State Hospital, Minsk, Belarus and Dorot. Medical Center for Rehabilitation and Geriatrics, Netanya, Israel.

Received: June 01, 2023; **Published:** June 14, 2023

Abstract

A 72-year-old man had with complaints of a debilitating cough, a change in voice, and a feeling of a foreign body in the throat for 4 months.

Keywords: *Lower Esophageal Sphincter (LES); Upper Esophageal Sphincter (UES); Gastroesophageal Reflux Disease*

Introduction and Case Report

A 72-year-old man had with complaints of a debilitating cough, a change in voice, and a feeling of a foreign body in the throat for 4 months. Very rarely, small pieces of food appear in the mouth. Within a month, he wakes up at night, as he chokes on saliva. He does not feel any acid or bitterness in his mouth. He has not heartburn, pain, or dysphagia. He noted the symptoms in detail in the questionnaire (Table 1). About 15 years ago he had heartburn and chest pressure. PPI treatment did not bring relief. The symptoms disappeared immediately after he swallowed contrast tablet with diameter 3 cm. Since then, he felt healthy, did not go to the doctors, did not take any treatment, and did not follow any dietary restrictions. In the table 1, the patient rated the severity of symptoms before treatment with a black circle and 25 days later with a red one.

X-ray examination of the esophagus, upper (UES) and lower esophageal sphincter (LES)

This method is based on the physiology of the esophagus and the esophageal sphincters. It is known that a significant increase in gastric pressure causes a reduction in both UES and LES [1]. Secondly, an increase in the tone of the LES causes a slowdown in the rate of esophageal peristalsis [2].

There is no restriction on pre-study meals. The patient lies on the X-ray table. Near his head is a jar with 200 ml of barium suspension. The patient drinks barium without interruption, and when the barium runs out he lifts his straightened legs high. At this point, the first radiograph is taken (Figure 1). The patient gets up and lies down again after 5 minutes. At this time, a second radiograph is taken at rest (Figure 2).

How concerned are you with the symptoms described below? 9.08.22 There is no symptom - 0. A strong symptom is 5. 4.09.22						
Hoarseness or alteration of voice	0	1	2	3	4	5
Sore throat or a desire to get rid of the irritant in the throat	0	1	2	3	4	5
Abundant sputum secreted from the back of the nose, or runny nose.	0	1	2	3	4	5
Difficulty in swallowing food, liquids or tablets	0	1	2	3	4	5
Cough after drinking, eating or resting in the horizontal position	0	1	2	3	4	5
Dyspnea or cases of sudden asphyxiation	0	1	2	3	4	5
Importunate cough	0	1	2	3	4	5
Sensation of foreign body in throat	0	1	2	3	4	5
Heartburn, chest pain or sensation of acid in the throat or mouth	0	1	2	3	4	5
Reflux Index (RI) $\Sigma = 23$	7					

Table 1: Questionnaire for the screening of gastroesophageal reflux disease.




Figure 1: High pressure in the stomach led to a reflex contraction of the UES and LES. Tight filling of the esophagus between the UES and LES is determined. LES is significantly shorter than normal (1.7 cm versus 3.6 cm). The phrenic ampulla (red line) is sharply dilated (4.4 cm versus 1.5 cm in normal). The walls of the esophagus are uneven with asymmetric waviness. At level D-4, a symmetrical constriction 6 mm wide with smooth contours is determined (arrow). Barium lodged in the hypopharynx is visible at the very top of the x-ray. The patient swallowed a 1.9 cm tablet, which passed into the stomach immediately, which confirmed the absence of stenosis. Spasm of the esophagus at the border of the upper and middle thirds indicates a high tone, i.e. high intraluminal pressure.



Figure 2: After 5 minutes at the rest, a spontaneous reflux of barium from the stomach into the esophagus is seen with a wide opening of the EGJ. The proximal sphincter, which most often closes the ampulla proximally, does not function in this case (on both images). Uneven longitudinal folds visible in the esophagus. But the barium didn't go above D-4 where there was constriction in the first radiograph. I believe that this is a functional sphincter that prevents acid from refluxing into the proximal esophagus. At this x-ray also seen barium stuck in the hypopharynx.

Discussion

The patient has a very weak LES. This follows firstly because it contracted as the pressure in the stomach rises, while in healthy people the force of the last peristaltic wave of the esophagus overcomes the tone of the LES and the bolus enters the stomach without delay. Therefore, in healthy individuals, LES does not contract and is not visible on the radiograph. Secondly, the length of the LES (1.7 cm) is two times shorter than the norm (3.6 cm), since its abdominal part opens in the stomach [3,4]. Thirdly, free reflux is visible in the second picture (Figure 2) in a horizontal position without pressure provocation. Therefore, he has episodes of reflux into the esophagus at night if the presence of chyme in the stomach. In milder cases, during reflux, the barium is retained in the ampulla, which is closed proximally by the functional sphincter. But this functional sphincter does not function in this case, but there is a functional narrowing at the level of the anatomical narrowing of the esophagus, due to the pressure of the aortic arch.

Probably at night, when the patient had swallowing saliva, it accumulated between this functional sphincter at the level of the aorta (arrow in figure 1) and the upper esophageal sphincter. When the esophagus filled with saliva between sphincters, resulted in increasing pressure, the upper esophageal sphincter cannot handle the pressure and relaxes. This causes saliva to reflux into the throat. After the passage of the tablet through this functional sphincter, its tone decreased, which contributes to the free passage of saliva through this zone:

1. This observation confirms numerous reports that GERD can progress for a long time without noticeable symptoms. From the current definition of GERD, it follows that if a person has no significant complaints, then he does not have GERD. Obviously, this definition is wrong! GERD is a chronic relapsing disease. It often progresses without clinical manifestations. The earlier the diagnosis is made and pathogenetic treatment is undertaken, the disease is easier, and the less serious complications occur.

2. The above x-ray examination lasted 10 minutes, during which two x-rays were taken. Its efficiency significantly exceeds the generally accepted method of X-ray examination. Tracking the passage of barium through the esophagus has low reliability in the diagnosis of GERD. Water-siphon test significantly increases the accuracy of diagnosis. However, it is not possible to evaluate it in numerical terms, since in all articles it was compared with pH monitoring, which was considered the gold standard without reason.

Conclusion

GERD with severe esophagitis. Pharyngitis.

Recommendations:

1. Consultation with an otolaryngologist.
2. Tab. Esomeprazole 20 mg x 2 (4 weeks).
3. Tab. Esomeprazole 20 mg x 1 (4 weeks).
4. Susp. Antacid 1 tablespoon 30 minutes after eating.
5. Stop eating food containing lactose.
6. Go to bed only with an empty stomach. Eat dinner 4-5 hours before bed.
7. Reduce the amount of food intake. Do not sit down for 30 minutes after eating.

The patient reported by phone that from the first night after taking the tablet, he did not get up at night and did not choke on saliva as before. The otolaryngologist found an inflammatory process in his pharynx and upper esophagus and prescribed an anti-inflammatory spray. The results of treatment 25 days after the examination are shown in table 1 in red. Reflux Index decreased from 23 to 7.

Bibliography

1. Shafik A., *et al.* "The effect of gastric overfilling on the pharyngo-esophageal and lower esophageal sphincter: a possible factor in restricting food intake". *Medical Science Monitor* 13.10 (2007): BR220-BR224.
2. Levin MD. GERD. Part II. Pathophysiology.
3. Chandrasoma P and DeMeester T. "A New Pathologic Assessment of Gastroesophageal Reflux Disease: The Squamo-Oxyntic Gap". *Advances in Experimental Medicine and Biology* 908 (2016): 41-78.
4. Levin MD. "Reaction to articles on high resolution manometry, the length of the lower esophageal sphincter and the diagnosis of gastroesophageal reflux disease". *Arquivos de Gastroenterologia* 56.2 (2019): 209-210.

Volume 6 Issue 7 July 2023

©All rights reserved by Michael D Levin.