

Esthetic Rehabilitation of a Discolored Maxillary Canine Tooth with a Porcelain Laminate Veneer

Arda Özdiler^{1*}, Hakan Bilhan² and Halit Yagci³

¹Researcher, Department of Prosthodontics, Istanbul University, Istanbul, Turkey

²Guest Professor, Department of Prosthodontics, Okan University, Istanbul, Turkey and Lecturer, Department of Prosthodontics, Witten/Herdecke University, Witten, Germany

³Private Practitioner, Portek-Dent Dental Laboratory, Bursa, Turkey

***Corresponding Author:** Arda Özdiler, Researcher, Department of Prosthodontics, Istanbul University, Istanbul, Turkey.

Received: August 25, 2018; **Published:** October 11, 2018

Abstract

Esthetic rehabilitation of a discolored single tooth in the esthetic zone presents major challenges for the clinicians. Due to the advances in dental materials, various conservative and minimal invasive treatment options are available in such cases. The application of the most appropriate treatment option is very important in terms of meeting the aesthetic demands of the patients. Thus, clinical success of a rehabilitation depends on the exact identification of a case and the decision of the appropriate treatment option. This article narrates the esthetic rehabilitation of a maxillary canine tooth due to fluorosis discoloration with a porcelain laminate veneer restoration.

Keywords: Esthetic Rehabilitation; Maxillary Canine Tooth; Porcelain Laminate Veneer

Introduction

Porcelain laminate veneers are one of the most conservative techniques applied in order to improve or reestablish the dental aesthetics [1-4]. These veneers are better in longevity and the resistance than the other veneer materials [5,6]. Porcelain laminate veneers can be applied in various cases such as discoloration, surface defects, diastema and malformation [1]. The current case report narrates the esthetic rehabilitation of a maxillary canine tooth due to fluorosis discoloration with a porcelain laminate veneer.

Case Report

A 25-year-old female patient reported her discontent with a discoloration at her maxillary left canine tooth (Figure 1). On the clinical examination of the patient, fluorosis related discolorations were observed at the labial medium 2/3 and the incisal part of her left maxillary canine tooth. Radiographic and clinical examination did not reveal any caries or periapical pathological condition. Occlusion was in a class 1 relationship, oral hygiene was good and the gingival tissues were healthy. As a result of the examinations, all appropriate treatment options for this case were described to the patient. Cad/Cam manufactured and individualized by cut-back method porcelain laminate veneer was accepted by the patient as a treatment option.



Figure 1: Discolorations at left maxillary canine tooth.

At the next clinical appointment tooth preparation was performed within minimally invasive approach followed by the final shade selection by utilizing a chromascop shade guide (Vita Toothguide 3D-Master; Vita, Zahnfabrik, Bad Sackingen, Germany). Final impressions were made using a polyvinyl-siloxane impression material (Elite-HD, Zhermack, Badia Polesine, Italy) by Putty-Wash technique. Temporary restoration was prepared with a self-curing composite material (Structur Premium, A2, VOCO, Cuxhaven, Germany) and cemented on the tooth surface by applying single point bonding technique (Single Bond, 3M ESPE, St Paul MN, USA).

Porcelain laminate veneer was manufactured by Cad/Cam method (Sirona X5 Scanner - Sirona Inlab 3D Software - Sirona MCX5 Milling Machine, Sirona Dental Systems, Germany; Vitablocs Triluxe Color 1M2, Vita Zahnfabrik, Bad Sackingen, Germany). Individualizations were made in cut-back technique to achieve the proper esthetic result (Vita VM 9 Porcelain, Vita Zahnfabrik, Bad Sackingen, Germany). The coherence and the fit of the laminate veneer were verified both individually and collectively on the prepared tooth. The patient was satisfied with the form, shade and adaptation of the restoration. Final cementation was done with a dual-cure resin cement (Variolink Esthetic Neutral, Ivoclar-Vivadent, Schaan, Lichtenstein) (Figure 2). The occlusion and contact correction with final polishing and finishing process was applied after 24 hours of cementation. Post-operative photos were taken and a night guard oral appliance was performed in order to prevent any trauma on the veneer caused by the parafunctional habits (Figure 3). One year post-operative evaluation did not reveal any fracture or discoloration of the veneer restoration (Figure 4). Patient was pleased with the conformity and esthetics of the restoration.



Figure 2: Clinical view after cementation process.



Figure 3: The occlusion and contact correction with final polishing and finishing process was applied after 24 hours of cementation.



Figure 4: One year post-operative view.

Discussion

Dental fluorosis is a common disorder affecting the enamel formation at the time of mineralization process. The clinical view exhibits a spectrum of change. Lusteless white lines or diffuse opacities are present in its mild form, while in the more severe forms generalized opaque and chalky appearance with confluent pitting and staining of hypo mineralized tissue may be seen [7]. Fluorosis distrups enamel significantly and affects appereance quite adversely which can cause adverse psychological effects on the individual. The treatment of enamel fluorosis usually ranges from ceramic veneer to direct bonding restorations and bleaching [8]. Although vital bleaching does improve the esthetics to certain extent it has only met with partial success in regard to moderate to severe fluorosis [9].

Porcelain veneers provide precise color match and translucency to the natural tooth and fulfill the need for adequate retention [10]. Veneers is considered to be a more conservative treatment approach compared to full-crown restorations because much less structure is removed from the tooth during the preparation of the veneers. For these types of restorations, the preparation does not cause the reduction of axial walls resulting in preservation of tooth structure and the surrounding hard and soft tissue architecture [11]. Thus, porcelain veneers are an appropriate type of restoration for young adults who have large pulp chambers close to the enamel. Preference of porcelain veneers to re-establish the esthetics in the anterior zone is considered as a satisfactory treatment option. Years of experience with both the technique and the materials utilized offer satisfactory, predictable, and lasting results [12]. Porcelain veneers restore the mechanical behavior and microstructure of the intact tooth *in vitro* even when they are bonded to an extensive dentin surface using an optimized application mode of dentine adhesives [13].

Conclusion

Veneering a discolored tooth is one of the most appropriate treatment options in the anterior section. Advances in the manufacturing techniques, structural properties of ceramics and luting cements, made porcelain veneering the most acceptable treatment option for the esthetic correction of anterior teeth than the other treatment options. However, clinical process requires high attention to all details in preparation and application process. In addition, adhesion, polishing and occlusion corrections are crucial for clinical success. Current case report presents the esthetic rehabilitation of a discolored maxillary canine tooth with a porcelain laminate veneer.

Bibliography

1. Stappert CF, *et al.* "Survival rate and fracture strength of maxillary incisors, restored with different kinds of full veneers". *Journal of Oral Rehabilitation* 32.4 (2005): 266-272.
2. Gresnigt M and Özcan M. "Esthetic rehabilitation of anterior teeth with porcelain laminates and sectional veneers". *Journal of the Canadian Dental Association* 77 (2011): 138-143.
3. Choi YS and Cho IH. "An effect of immediate dentin sealing on the shear bond strength of resin cement to porcelain restoration". *Journal of Advanced Prosthodontics* 2.2 (2010): 39-45.
4. Peumans M, *et al.* "A prospective ten year clinical trial of porcelain veneers". *Journal of Adhesive Dentistry* 6.1 (2004): 65-76.
5. Wiedhahn K, *et al.* "Clinical long term results with 617 Cerec veneers: A nine-year report". *International Journal of Computerized Dentistry* 8.3 (2005): 233-246.
6. Kelly JR, *et al.* "Ceramic in dentistry: History and historical roots and current perspectives". *Journal of Prosthetic Dentistry* 75.1 (1996): 18-32.
7. Fejerskov O, *et al.* "The ultrastructure of fluorosed human dental enamel". *European Journal of Oral Sciences* 82.5 (1974): 357-372.
8. Nazirkar G and Meshram S. "An Evaluation of Two Modern All-Ceramic Crowns and their comparison with Metal Ceramic Crowns in terms of the Translucency and Fracture Strength". *International Journal of Dental Clinics* 3.1 (2011): 5-7.
9. Sherwood IA. "Fluorosis varied treatment options". *Journal of Conservative Dentistry* 13.1 (2010): 47-53.
10. Ishikawa-Nagai S, *et al.* "Clinical evaluation of perceptibility of color differences between natural teeth and all-ceramic crowns". *Journal of Dentistry* 37.1 (2009): e57-e63.
11. Chen YW and Raigrodski AJ. "A conservative approach for treating young adult patients with porcelain laminate veneers". *Journal of Esthetic and Restorative Dentistry* 20.4 (2008): 223-236.
12. McLean JW. "Evolution of dental ceramics in the twentieth century". *Journal of Prosthetic Dentistry* 85.1 (2001): 61-66.
13. Magne P and Douglas WH. "Porcelain veneers: Dentin bonding optimization and biomimetic recovery of the crown". *International Journal of Prosthodontics* 12.2 (1999): 111-121.

Volume 17 Issue 11 November 2018

©All rights reserved by Arda Özdiler., *et al.*