

Coagulation, Fibrinolytic and Cytokines Parameters Response to Weight Reduction in Obese Subjects

Shehab Mahmoud Abd El-Kader^{1*} and Osama H Al-Jiffri²

¹Department of Physical Therapy, Faculty of Applied Medical Sciences, King Abdulaziz University, Saudi Arabia

²Department of Medical Laboratory Technology, Faculty of Applied Medical Sciences, King Abdulaziz University, Saudi Arabia

***Corresponding Author:** Shehab Mahmoud Abd El-Kader, Department of Physical Therapy, Faculty of Applied Medical Sciences, King Abdulaziz University, Jeddah, Saudi Arabia.

Received: December 30, 2016 **Published:** January 09, 2017

Abstract

Background: Obesity is a major risk factor for developing cardiovascular disorders due to increased platelets count and platelets activation. Objective: The aim of this study was to measure the influence of weight loss on the coagulation and fibrinolysis parameters among obese patients. Material and Methods: One hundred obese subjects of both genders (mean age 46.72 ± 3.13 year) were selected from the Internal Medicine Department at King Abdul Aziz University Teaching Hospital, their body mass index (BMI) ranged from 32 to 36 Kg/m². Participants were enrolled into two study groups; the first group (A) received aerobic exercises and diet regimen for 12 weeks, the second group (B) received no training intervention. Results: Results of group (A) showed a statistical significant decrease in the mean values of von Willbrand factor antigen (vWF-Ag), fibrinogen, plasminogen activator inhibitor-1 antigen (PAI-1:Ag) and plasminogen activator inhibitor-1 activity (PAI-1:Ac), tumor necrotic factor-alpha (TNF- α), Interleukin - 6 (IL-6) and body mass index (BMI), while the mean values of partial thromboplastin time (PTT), prothrombin time (PT), tissue plasminogen activator antigen (tPA:Ag) and tissue plasminogen activator activity (tPA:Ac) increased significantly after 3 months of the weight reduction program. However, the mean values of all investigated parameters of group (B) had no significant changes. Moreover, comparison between both groups had statistical significant differences in all investigated parameters at the end of the study ($P < 0.05$). Conclusion: weight reduction program modulates systemic inflammation, fibrinolytic and coagulation of obese subjects.

Keywords: Obesity; Inflammatory Cytokines; Coagulation; Fibrinolytic Parameters

Introduction

Obesity is a major risk factor for developing cardiovascular disorders due to increased platelets count and platelets activation [1-3], disturbed blood lipid profile, endothelial dysfunction systemic inflammation and atherosclerosis [4,5]. Despite all the improved knowledge on the pathogenesis and treatment of atherothrombosis associated with obesity, it is predicted that coronary heart disease will be the dominant cause of mortality worldwide by 2020 [6].

Obesity is usually associated with reduction of fat tissue oxygenation leads to production of inflammatory cytokines [7] as tumor necrosis factor- α , interleukin-6 [8], leptin and resistin produced as a result of adipose tissue dysfunction [9]. Increased levels of systemic inflammation markers as Interleukin-6 (IL-6) which induces increased platelet count size and function [10]. Moreover, increased biomarkers of platelet adhesion and endothelial cell function can be induced by the high level of serum of tumor necrosis factor- alpha (TNF- α) [11].

Regular exercise training can modulate the coagulation and fibrinolytic parameters [12] because of reduction of body fat tissue, which may reduce the cardiovascular system disorders (CVD) [13-15]. Also, weight reduction can have a great contribution in limitation of morbidity and mortality related to CVD through amelioration of endothelial dysfunction and fibrinolytic abnormalities [16,17].

Citation: Shehab Mahmoud Abd El-Kader and Osama H Al-Jiffri. "Coagulation, Fibrinolytic and Cytokines Parameters Response to Weight Reduction in Obese Subjects". *EC Gastroenterology and Digestive System* 1.5 (2017): 179-185.

The aim of this study was to measure the influence of weight loss on the coagulation and fibrinolysis parameters among obese patients.

Materials and Methods

Subjects

One hundred obese subjects of both genders (mean age 46.72 ± 3.13 year) were selected from the Internal Medicine Department at King Abdul Aziz University Teaching Hospital, their body mass index (BMI) ranged from 32 to 36 Kg/m². Smoking, hypertension, diabetes, hyperthyroidism, cardiovascular disorders and drugs affecting coagulation were the exclusion criteria. Participants were enrolled into 2 study groups; the first group (A) received aerobic exercises and diet regimen for 12 weeks, the second group (B) received no training intervention. All participants signed the consent form before sharing in the study.

Methods

Equipment

- 1) Treadmill (Enraf Nonium, Model display panel Standard, NR 1475.801, Holland) was used in performance of aerobic walking exercise.
- 2) Weight and height scale (JENIX DS 102, Dongsang, South Korea) was used to measure weight and height to calculate the body mass index (BMI).

Measurements

Laboratory analysis: Overnight venous blood samples were drawn, centrifuged and stored frozen at -80°C for analysis of plasma Willbrand factor antigen (vWF-Ag), fibrinogen, plasminogen activator inhibitor-1 antigen (PAI-1:Ag), plasminogen activator inhibitor-1 activity (PAI-1:Ac), partial thromboplastin time (PTT), prothrombin time (PT), tissue plasminogen activator antigen (tPA:Ag) and tissue plasminogen activator activity (tPA:Ac) employing the ST-4 coagulation instrument (Zymutest Fibrinogen, ELISA, Hyphen Biomed, Neuville sur Oise, France). Tumor necrotic factor - alpha (TNF- α) and Interleukin - 6 (IL-6) levels were analyzed with ELISA kits with ELISA microplate strip washer in addition to ELISA microplate reader.

Evaluation of anthropometric parameters: Body weight was measured with a calibrated balance scale (HC4211, Cas Korea, South Korea) and the body height was measured using (JENIX DS 102, Dongsang, South Korea), while body mass index (BMI) was determined by equation: $\text{BMI} = \text{Body weight} / (\text{Height})^2$.

All measurements of BMI, vWF, PT, PTT, PAI-1, TPA, fibrinogen TNF- α and IL-6 were taken before the study (pre-test) and after three months (post-test).

Procedures

Following the previous evaluation, all participants were included into two study groups:

Study group (A): Participants practiced treadmill aerobic exercise training. Training program included 5 minutes for warming-up in the form of range motion and stretching exercises, 30 minutes of aerobic exercise training with intensity equal 60 - 70% of the individual maximum heart rate followed by cooling down for 10 minutes (on treadmill with low speed and without inclination). Participants had 3 sessions /week for 3 months with close supervision of physical therapist. In addition, a dietician performed an interview-based food survey for all participants of group (A) for detection of feeding habits, abnormal dietary behavior and to prescribe the balanced low caloric diet that provided 1200 Kilocalories/day for 12 weeks. The same dietitian continuously monitored all participant caloric intake through reviewing the detailed record of food intake every 2 weeks by the dietitian.

Group (B): Participants only received their usual medical therapy for 3 months.

Statistical Analysis

Mean values of the investigated parameters will be compared by student paired "t" test. While, the unpaired" test will be used to compare between the two groups ($P < 0.05$).

Results

Results of group (A) showed a statistical significant decrease in the mean values of von Willbrand factor antigen (vWF-Ag), fibrinogen, plasminogen activator inhibitor-1 antigen (PAI-1:Ag) and plasminogen activator inhibitor-1 activity (PAI-1:Ac), tumor necrotic factor - alpha (TNF- α), Interleukin-6 (IL-6) and body mass index (BMI), while the mean values of partial thromboplastin time (PTT), prothrombin time (PT), tissue plasminogen activator antigen (tPA:Ag) and tissue plasminogen activator activity (tPA:Ac) increased significantly after 3 months of the weight reduction program. However, the mean values of all investigated parameters of group (B) had no significant changes (Table 1 and 2). Moreover, comparison between both groups had statistical significant differences in all investigated parameters at the end of the study (Table 3) ($P < 0.05$).

	Mean \pm SD		t- value	P-value
	Pre	Post		
BMI (Kg / m ²)	35.63 \pm 5.71	26.92 \pm 5.43*	8.21	0.002
PTT (s)	20.53 \pm 3.46	24.72 \pm 3.38*	6.17	0.008
PT (s)	10.62 \pm 1.74	12.81 \pm 1.65*	5.36	0.006
Fibrinogen (mg/mL)	3.91 \pm 0.83	2.41 \pm 0.82*	4.23	0.017
tPA:Ac (IU/mL)	5.27 \pm 0.94	6.83 \pm 0.86*	4.75	0.013
tPA:Ag (ng/mL)	3.13 \pm 0.72	4.82 \pm 0.68*	4.31	0.018
vWF-Ag (%)	91.26 \pm 11.27	72.82 \pm 10.88*	9.18	0.001
PAI-1:Ag (ng/mL)	18.26 \pm 3.91	10.12 \pm 3.78*	5.42	0.005
PAI-1:Ac (AU/mL)	4.73 \pm 0.86	3.24 \pm 0.75*	4.63	0.019
IL-6 (pg/mL)	8.51 \pm 2.16	5.75 \pm 1.98*	5.22	0.003
TNF- α (pg/mL)	5.43 \pm 1.43	4.11 \pm 1.25*	4.63	0.012

Table 1: Mean value and significance of the pre and posttest values of PT, BMI, PTT, tPA:Ag, tPA:Ac, vWF-Ag, fibrinogen, PAI-1:Ag, PAI-1:Ac, TNF- α and IL-6 of the training group.

BMI: Body Mass Index; PT: Prothrombin Time; PTT: Partial Thromboplastin Time; tPA:Ac: Tissue Plasminogen Activator Activity; tPA:Ag: Tissue Plasminogen Activator Antigen; vWF-Ag: Von Willbrand Factor Antigen; PAI-1:Ac: Plasminogen Activator Inhibitor-1 Activity; PAI-1:Ag: Plasminogen Activator Inhibitor-1 Antigen; TNF-A: Tumor Necrotic Factor - Alpha; IL-6: Interleukin- 6; (*): Indicates A Significant Difference, $P < 0.05$.

	Mean \pm SD		t- value	P-value
	Pre	Post		
BMI (Kg / m ²)	35.73 \pm 5.61	34.28 \pm 5.47	1.12	1.453
PTT (s)	20.82 \pm 3.21	21.12 \pm 3.20	0.74	1.976
PT (s)	11.15 \pm 1.82	11.63 \pm 1.85	0.81	1.438
Fibrinogen (mg/mL)	3.88 \pm 0.84	3.79 \pm 0.81	0.64	1.876
tPA:Ac (IU/mL)	5.62 \pm 0.74	5.48 \pm 0.69	0.68	1.641
tPA:Ag (ng/mL)	3.22 \pm 0.59	3.17 \pm 0.53	0.79	1.912

vWF-Ag (%)	92.15 ± 11.23	90.14 ± 11.18	1.25	1.823
PAI-1:Ag (ng/mL)	18.74 ± 3.81	18.16 ± 3.42	0.51	1.754
PAI-1:Ac (AU/mL)	4.83 ± 0.91	4.75 ± 0.86	0.66	1.883
IL-6 (pg/mL)	8.32 ± 2.24	8.18 ± 2.21	0.62	1.741
TNF-α (pg/mL)	5.49 ± 1.37	5.31 ± 1.35	0.65	1.864

Table 2: Mean value and significance of the pre and posttest values of PT, BMI, PTT, tPA:Ag, tPA:Ac, vWF-Ag, fibrinogen, PAI-1:Ag, PAI-1:Ac, TNF-α and IL-6 of the control group.

BMI: Body Mass Index; PT: Prothrombin Time; PTT: Partial Thromboplastin Time; tPA:Ac: Tissue Plasminogen Activator Activity; tPA:Ag: Tissue Plasminogen Activator Antigen; vWF-Ag: Von Willbrand Factor Antigen; PAI-1:Ac: Plasminogen Activator Inhibitor-1 Activity; PAI-1:Ag: Plasminogen Activator Inhibitor-1 Antigen; TNF-A: Tumor Necrotic Factor - Alpha; IL-6: Interleukin- 6.

	Mean ± SD		t- value	P-value
	Training group	Control group		
BMI (Kg / m ²)	26.92 ± 5.43	34.28 ± 5.47*	6.23	0.012
PTT (s)	24.72 ± 3.38	21.12 ± 3.20*	4.82	0.023
PT (s)	12.81 ± 1.65	11.63 ± 1.85*	4.21	0.018
Fibrinogen (mg/mL)	2.41 ± 0.82	3.79 ± 0.81*	3.72	0.011
tPA:Ac (IU/mL)	6.83 ± 0.86	5.48 ± 0.69*	3.64	0.012
tPA:Ag (ng/mL)	4.82 ± 0.68	3.17 ± 0.53*	3.51	0.016
vWF-Ag (%)	72.82 ± 10.88	90.14 ± 11.18*	7.33	0.013
PAI-1:Ag (ng/mL)	10.12 ± 3.78	18.16 ± 3.42*	4.22	0.028
PAI-1:Ac (AU/mL)	3.24 ± 0.75	4.75 ± 0.86*	3.61	0.027
IL-6 (pg/mL)	5.75 ± 1.98	8.18 ± 2.21*	4.23	0.015
TNF-α (pg/mL)	4.11 ± 1.25	5.31 ± 1.35*	4.16	0.019

Table 3: Mean value and significance of the posttest values PT, BMI, PTT, tPA:Ag, tPA:Ac, vWF-Ag, fibrinogen, PAI-1:Ag, PAI-1:Ac, TNF-α and IL-6 of the training and control groups.

BMI: Body Mass Index; PT: Prothrombin Time; PTT: Partial Thromboplastin Time; tPA:Ac: Tissue Plasminogen Activator Activity; tPA:Ag: Tissue Plasminogen Activator Antigen; vWF-Ag: Von Willbrand Factor Antigen; PAI-1:Ac: Plasminogen Activator Inhibitor-1 Activity; PAI-1:Ag: Plasminogen Activator Inhibitor-1 Antigen; TNF-A: Tumor Necrotic Factor - Alpha; IL-6: Interleukin- 6; (*): Indicates A Significant Difference, P < 0.05.

Discussion

Disorders of coagulation and fibrinolysis have a great contribution in obesity related cardiovascular disorders [18-19]. We found that weight loss program improves the parameters of systemic inflammation along with amelioration of coagulation disorders in obese subjects, our findings were in line with many previous studies.

Barbeau and colleagues stated that weight loss can reduce the cardiovascular disorders (CVD) mortality rate via modulation of endothelial and fibrinolytic dysfunction [20]. Svendsen, *et al.* proved that weight reduction can control oxidative stress, platelets activation and chronic inflammation in obese women [21]. In addition, Anfossi and colleagues found that 10% body weight loss is able to correct platelet abnormalities related to central obesity related insulin resistance [1].

Concerning PAI-1 activity & antigen and tPA activity & tPA antigen our results showed that weight loss significantly increase the values of tPA antigen and tPA activity and reduce the values of PAI-1 activity and antigen, these findings are in line with many previous trials of weight loss [20-22]. In addition, Murakami and colleagues stated that increased values of PAI-1 activity were positively correlated with high values of BMI in obese subjects, which were reduced as a result to weight loss [13].

Regarding PTT, PT and vWF-Ag, our results revealed an increase in the mean values of PTT and PT because of weight loss and these findings agreed with Piccone, *et al.* [23] and Stratton, *et al.* concluded that a six months of endurance exercises resulted in reduction in fibrinogen and PAI-1 activity and enhanced the t-PA activity in elderly men [24]. However, there was a significant reduction of vWF and these results are in line with Saenko, *et al.* [25] and Paton, *et al.* stated that serum vWf:Ag and tPA antigen were reduced in ten subjects performed two maximal oxygen uptake (VO_{2max}) test trials [26].

Concerning our results regarding the serum level fibrinogen, weight loss led to reduction in fibrinogen concentration, this finding is in line with De Souza, *et al.* proved that physically active postmenopausal women have lower serum fibrinogen, t-PA antigen, PAI-1 antigen, and PAI-1 activity and higher ($P < 0.01$) t-PA activity levels than sedentary postmenopausal women [12], which may be due to reduced body fat mass as a result of regular exercises [27,28].

The results of this study indicated a great significant reduction in TNF- α , IL6 and BMI only in-group (A) this means that weight reduction is an effective anti-inflammatory treatment, these results agreed with Esposito, *et al.* proved that weight reduction led to reduction in inflammatory cytokines and increase the serum levels of anti-inflammatory cytokines in obese women [29]. Reinher and colleagues clearly stated that life style modification program reduces serum level of TNF- α [30]. Also, Gallistat, *et al.* stated that a three weeks weight reducing program resulted in significant decrease in IL-6, body fat mass and BMI [31].

Conclusion

Weight reduction program modulates systemic inflammation, fibrinolytic and coagulation of obese subjects.

Acknowledgment

This project was funded by the Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah, under grant no. (57-142-1434). The authors, therefore, acknowledge with thanks DSR technical and financial support.

Bibliography

1. Anfossi G., *et al.* "Platelet dysfunction in central obesity". *Nutrition, Metabolism and Cardiovascular Diseases* 19.6 (2009): 440-449.
2. Després J. "Cardiovascular disease under the influence of excess visceral fat". *Critical Pathways in Cardiology* 6.2 (2007): 51-59.
3. Samocha-Bonet D., *et al.* "Platelet counts and platelet activation markers in obese subjects". *Mediators of Inflammation* (2008): 834153.
4. Ritchie S and Connell J. "The link between abdominal obesity, metabolic syndrome and cardiovascular disease". *Nutrition, Metabolism and Cardiovascular Diseases* 17.4 (2007): 319-326.
5. Galic S., *et al.* "Adipose tissue as an endocrine organ". *Molecular and Cellular Endocrinology* 316.2 (2010): 129-139.
6. Vilahur G and Badimon L. "Antiplatelet properties of natural products". *Vascular Pharmacology* 59.3-4 (2013): 67-75.
7. Hulsmans M and Holvoet P. "The vicious circle between oxidative stress and inflammation in atherosclerosis". *Journal of Cellular and Molecular Medicine* 14.1-2 (2010): 70-78.

8. Surmi B and Hasty A. "The role of chemokines in recruitment of immune cells to the artery wall and adipose tissue". *Vascular Pharmacology* 52.1-2 (2010): 27-36.
9. Vasconcelos E., *et al.* "Reactive oxygen species generation in peripheral blood monocytes and oxidized LDL are increased in hyperlipidemic patients". *Clinical Biochemistry* 42.12 (2009): 1222-1227.
10. Peng J., *et al.* "Relative reactivity of platelets from thrombopoietin- and interleukin-6-treated dogs". *Blood* 87.10 (1996): 4158-4163.
11. Lou J., *et al.* "Platelets play an important role in TNF-induced microvascular endothelial cell pathology". *American Journal of Pathology* 151.5 (1997): 1397-1405.
12. DeSouza C., *et al.* "Physical activity status and adverse age-related differences in coagulation and fibrinolytic factors in women". *Arteriosclerosis, Thrombosis, and Vascular Biology* 18.3 (1998): 362-368.
13. Murakami T., *et al.* "Impact of weight reduction on production of platelet-derived microparticles and fibrinolytic parameters in obesity". *Thrombosis Research* 119.1 (2007): 45-53.
14. Jahangard T., *et al.* "The effect of short-term aerobic training on coagulation and fibrinolytic factors in sedentary healthy postmenopausal women". *Maturitas* 64.4 (2009): 223-227.
15. Ades P., *et al.* "The influence of obesity and consequent insulin resistance on coronary risk factors in medically treated patients with coronary disease". *International Journal of Obesity* 32.6 (2008): 967-974.
16. Hamdy O., *et al.* "Lifestyle modification improves endothelial function in obese subjects with the insulin resistance syndrome". *Diabetes Care* 26.7 (2003): 2119-2125.
17. National Institutes of Health. "Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults—the evidence report". *Obesity Research* 6.2 (1998): 51S-209S.
18. Lee I., *et al.* "Physical activity and coronary heart disease in women: "is no pain, no gain" passé?" *Journal of the American Medical Association* 285.11 (2001): 1447-1454.
19. Szymanski L., *et al.* "Relationship of physical fitness, hormone replacement therapy, and hemostatic risk factors in postmenopausal women". *Journal of Applied Physiology* 98.4 (2005): 1341-1348.
20. Barbeau P., *et al.* "Hemostatic and inflammatory markers in obese youths: effects of exercise and adiposity". *Journal of Pediatrics* 141.3 (2002): 415-420.
21. Svendsen O., *et al.* "Plasminogen activator inhibitor-1, tissue-type plasminogen activator, and fibrinogen: effect of dieting with or without exercise in overweight postmenopausal women". *Arteriosclerosis, Thrombosis, and Vascular Biology* 16.3 (1996): 381-385.
22. Mavri A., *et al.* "Impact of adipose tissue on plasma plasminogen activator inhibitor-1 in dieting obese women". *Arteriosclerosis, Thrombosis, and Vascular Biology* 19.6 (1999): 1582-1587.
23. Piccone G., *et al.* "Exercise-induced change in clotting times and fibrinolytic activity during official 1600 and 2000 meters trot races in standard horses". *Acta Veterinaria Brunensis* 74 (2005): 509-514.

24. Stratton J., *et al.* "Effect of physical conditioning on fibrinolytic variables and fibrinogen in young and old healthy adults". *Circulation* 83.5 (1991): 1692-1697.
25. Saenko E., *et al.* "Role of low-density lipoprotein-related protein receptor in mediation of factor VIII catabolism". *Journal of Biological Chemistry* 274.53 (1999): 37685-37692.
26. Paton C., *et al.* "Changes in von Willebrand factor and fibrinolysis following a post-exercise cool-down". *European Journal of Applied Physiology* 92.3 (2004): 328-333.
27. Stefanick M., *et al.* "Distribution and correlates of plasma fibrinogen in middle-aged women: initial findings of the Postmenopausal Estrogen/Progestin Interventions (PEPI) study". *Arteriosclerosis, Thrombosis, and Vascular Biology* 15.12 (1995): 2085-2093.
28. Krobot K., *et al.* "Determinants of plasma fibrinogen: relation to body weight, waist-to-hip ratio, smoking, alcohol, age, and sex: results from the second MONICA Augsburg Survey 1989-1990". *Arteriosclerosis, Thrombosis* 12.7 (1992): 780-788.
29. Esposito K., *et al.* "Effect of weight loss and lifestyle changes on vascular inflammatory markers in obese women: a randomized trial". *Journal of the American Medical Association* 289.14 (2003): 1799-1804.
30. Reinher T., *et al.* "High-sensitive C-reactive protein, tumour necrosis factor α , and cardiovascular risk factors before and after weight loss in obese children". *Metabolism* 54.9 (2005): 1155-1161.
31. Gallistatl S., *et al.* "Changes in serum interleukin-6 concentrations in obese children and adolescents during a weight reduction program". *International Journal of Obesity* 25.11 (2001): 1640-1643.

Volume 1 Issue 5 January 2017

© All rights reserved by Shehab Mahmoud Abd El-Kader and Osama H Al-Jiffri.