

## An Eighteen Year Study of Intestinal Protozoans in the Los Angeles Area Between 1996 and 2013\*

Omar M Amin\* and Karim O Amin

Department of Parasitology, Parasitology Center, Inc. (PCI), 11445 E. Via Linda #2-419, Scottsdale, Arizona, United States

\*Corresponding Author: Omar M Amin, Department of Parasitology, Parasitology Center, Inc. (PCI), 11445 E. Via Linda #2-419, Scottsdale, Arizona, United States.

Received: April 22, 2017; Published: May 17, 2017

### Abstract

Seasonal and annual prevalence of intestinal protozoans were studied for a period of 18 years in an urban/suburban Los Angeles area. This is the first such study anywhere in the world. A total of 7766 fecal specimens from 3883 patients in the Los Angeles County, from 1996 to 2013 were tested at Parasitology Center, Inc. (PCI), Scottsdale, Arizona. During this period, 1629 (41%) of patients were found infected with one or more protozoan parasites. The most prevalent parasites were *Blastocystis hominis* (19%), *Entamoeba histolytica/E. dispar* (6%) *E. hartmanni* (6%), and *Cryptosporidium parvum* (5%). *Blastocystis hominis* made up 45% of all infections. Infections with *Blastocystis hominis* progressively declined through 2013 while those of *C. parvum* increased. Infections with *B. hominis* were more prevalent in colder weather and lowest in August and September. Infections with *C. parvum* were most prevalent from March to June and lowest in August. The overall monthly prevalence for all protozoan parasites varied between 34% in August and 51% in February. The composition of the parasitic fauna diagnosed, annual prevalence rates, and seasonality were discussed in comparison with other studies.

**Keywords:** Intestinal Protozoans; Los Angeles; 1996-2013; Prevalence; Seasonality

### Introduction

Parasitological studies of large patient populations are rare in the United States compared to third-world countries where endemic parasitosis are more readily reported [1]. We, At the Parasitology Center, Inc. (PCI), in Scottsdale, Arizona, routinely monitor and report on the patterns and trends of human parasitosis in the US. The seasonal prevalence of 19 species of intestinal parasites infecting 916 of 2,896 (32%) examined patients from 48 states in the year 2000 was reported [2]. In that study, 314 of 859 examined patients (36%) from California were infected [2]. Infections with helminth parasites such as *Ascaris lumbricoides* and non-major protozoans were rare and are not included in the present study. The seven reported species of protozoan parasites constituted 91.5% of 18 species of parasites reported in the United States. Multiple infections with 2-4 parasitic species constituted 10% of the infected cases [2]. We also researched the epidemiology of *Blastocystis hominis* in 48 states and the District of Columbia in 2002 - 2004 and included trends in annual, seasonal, geographical and host distribution, and symptomology by age, sex and season [3]. In that report, 16% of 10,582 fecal specimens from 5,291 patients tested positive for *B. hominis*; in California, 263 of 1,328 examined patients (20%) were also positive for *B. hominis*. In a similar 3-year epidemiological study of 9,856 fecal specimens from 4,928 patients from all states and the District of Columbia that we tested between 2003 and 2005, 279 (6%) were positive for *Cryptosporidium parvum* infections. Studies of this magnitude are not known in the US. Few other studies of relatively large patient populations in the US [4,5] or in more geographically limited populations addressing *B. hominis* only, e.g., California [6] or Ontario [7] have been reported. The present investigation is the first to cover the span of 18 years. Evaluating the patterns and trends of parasitic infections in studies of such a long duration is a great tool for understanding the epidemiological characteristics, disease burden, improving the reporting of cases, planning prevention, therapeutic, and other public health measures to

be considered in the management of these infections. Nevertheless, an overview of studies of shorter duration from comparable urban/suburban area in developing and developed nations are included.

## Materials and Methods

A total of 7766 specimens from 3883 patients (two specimens per patient) were collected, preserved, and transported to Parasitology Center, Inc. (PCI) in Proto-fix™ (Alpha-Tec Systems, Inc. Vancouver, Wash.) or SAF (sodium acetate–acetic acid–formalin mixture) in plastic vials provided in mailable kits. Patients were referred to PCI by 187 doctors in Los Angeles County from January 1996 through December 2013. The number of referring practitioners varied from year to year. Specimens were processed and stained with CONSED™ according to the manufacturer's (Alpha-Tec Systems) directions. This procedure was used in thousands of specimens at PCI, evaluated, and described previously [8]. Briefly, specimens are filtered, mixed with CONSED and ethyl acetate, vortexed, centrifuged, and decanted. The resulting fecal plug was mixed with CONSED diluting reagent, transferred to, and mounted on a slide for microscopic examination as wet mounts. All samples were evaluated by the same observer blinded to patient information. The reliability of diagnosis is indicated by the consistency of detection of different parasites at different levels of infection during the same period of time. Differences in the number of patient samples tested in different years reflect changing patterns of patient traffic from Los Angeles County over time. Positive results were quantified (number of organisms per high power field on a scale of 1 - 4) from duplicate samples from each patient. About 10% of infected patients had 2 - 4 parasitic species each which is the same prevalence rate of concurrent infections noted in our earlier study [2]. A prevalence rate based on the number of infected/examined patients would be about 10% lower but would not reflect the activity of individual parasitic species. The number of samples tested underwent a gradual decrease over the years corresponding to changing patterns of patients and practitioners' traffic in the Los Angeles area while increasing elsewhere in the US and internationally.

## Issues in Diagnosis

We accept the possible presence of *Blastocystis* and *Cryptosporidium* organisms from animal sources in human infections with *B. hominis* and *Cryptosporidium parvum* as reviewed in Tan [9] and Garcia [10], respectively. Fletcher, *et al.* [11] provided a comprehensive, well referenced diagnostic coverage of intestinal protozoan infections in developed countries. Human, mammalian, avian, and reptilian isolates of *B. hominis* have been assigned to 13 subtypes. *Blastocystis* subtype 3 is most commonly associated with illness in human prevalence studies. The term *Blastocystis hominis* normally refers to about 10 different genetic populations that are indistinguishable microscopically. That term is used for parasites isolated from humans while *Blastocystis* spp. is used for isolations from animal hosts [11]. Most animals are not infected with human pathogenic cryptosporidiosis. However, zoonotic transmission from direct contact with infected animals or their feces through indirect sources including drinking of contaminated water occurs. The invasive *Entamoeba histolytica* trophozoites is less common than the morphologically identical non-pathogenic *E. dispar* and *E. moshkovskii* but distinguishable from them by isoenzyme analysis. *Giardia intestinalis* infections are detected microscopically by us and also by various antigen assays demonstrating seven genetically distinct genotypes (A-G). Assemblages A and B infect humans as well as other mammalian species and are considered zoonotic. See Fletcher, *et al.* [11] for a discussion of above diagnostic issues.

## Results

### Prevalence

A total of 1629 parasitic infections from 3883 patients (41%) were identified between 1996 - 2013 in the Los Angeles area. *Blastocystis hominis* was the most frequent parasitological finding. It was identified in 19% of samples, and represented 45% of all parasitological infections (Table 1). The next most common protozoan was *E. histolytica/dispar*. It was found in 6% of samples constituting 15% of all infections. The prevalence of *E. hartmanni*, *C. parvum* and *E. coli* was 6%, 5%, and 4%, respectively. *Dientamoeba fragilis* and *Giardia intestinalis* were found in 4% and < 1% of the samples examined, in the same order. These parasites constituted 91.5% of 18 species of intestinal parasites reported from 5792 fecal specimens tested from throughout the US in 2000 [2].

Parasite species	Patients infected	Prevalence	Percent of infections
<i>Blastocystis hominis</i>	732	19%	45
<i>Entamoeba histolytica/E.dispar</i> **	234	6%	15
<i>Entamoeba hartmanni</i>	226	6%	14
<i>Cryptosporidium parvum</i>	201	5%	12
<i>Entamoeba coli</i> **	156	4%	10
<i>Dientamoeba fragilis</i>	60	1%	3
<i>Giardia intestinalis</i> **	20	0%***	1
Total	1629*	41%	100

**Table 1:** Prevalence of protozoan infections diagnosed from 3883 patients\* examined from Los Angeles County between 1996 and 2013.

\*About 10% of infected patients were concurrently infected with more than 1 species of parasite.

\*\*Trophozoites and cysts.

\*\*\*0% indicates values of less than 1%.

**Annual Prevalence**

The prevalence of all parasitic infections was highest during 1996 - 97 (63%) then gradually declined through 2012-2013. *Giardia intestinalis* was the only protozoan that was consistently identified in less than 1% of tested samples throughout the study period. The prevalence of *Cryptosporidium* sp. increased and decreased cyclically reaching the lowest level of 2% in 1996-1997 (when other protozoans were most prevalent) to 11% in 2008 - 2009 before declining again. The prevalence of *B. hominis* infections progressively declined from 21% in 1996-2001 to 7% in 2012 - 2013 (Table 2). The total number of samples submitted from Los Angeles County was highest in 1996 - 1997 but declined then stabilized afterwards as did the total prevalence rates of all parasites.

Combined 2 year periods	96-97	98-99	00-01	02-03	04-05	06-07	08-09	10-11	12-13
<i>Blastocystis hominis</i>	315(21)*	203(21)	112(21)	29(15)	26(14)	18(9)	12(8)	9(9)	8(7)
<i>Cryptosporidium parvum</i>	30(2)	72(8)	18(3)	14(7)	14(10)	16(8)	17(11)	9(9)	6(4)
<i>Dientamoeba fragilis</i>	47(3)	1(0)	1(0)	1(1)	0(0)	2(2)	1(1)	0(0)	5(5)
<i>Entamoeba coli</i> cysts	50(3)	17(2)	6(1)	3(2)	4(2)	3(1)	2(1)	1(1)	1(0)
<i>E. coli</i> trophozoites	62(4)	3(0)	4(1)	1(1)	1(1)	0(0)	0(0)	0(0)	0(0)
<i>E. hartmanni</i> cysts	89(6)	9(1)	0(0)	0(0)	1(1)	0(0)	0(0)	0(0)	1(0)
<i>E. hartmanni</i> trophozoites	57(4)	14(1)	4(1)	1(0)	0(0)	0(0)	0(0)	1(1)	0(0)
<i>E. histolytica/dispar</i> cysts	6(2)	10(3)	8(2)	2(1)	10(3)	10(3)	8(3)	4(1)	4(1)
<i>E. histolytica/dispar</i> trophozoites	145(10)	12(1)	1(0)	2(1)	0(0)	0(0)	0(0)	0(0)	0(0)
<i>Giardia intestinalis</i> cysts	9(1)	5(1)	2(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)
<i>G. intestinalis</i> trophozoites	1(0)**	1(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)
Total # of infections	927	337	138	51	47	39	32	20	21
Total examined	1483	947	530	193	187	195	155	92	101
Prevalence	63%	36%	26%	26%	25%	20%	21%	20%	21%

**Table 2:** Number and prevalence (%) of infections with major protozoan parasites diagnosed from 3883 patients examined from Los Angeles County between 1996 and 2013 by two year period increments.

\*No. of infections (% of examined patients per each 2 year period). Percentages are rounded

\*\*0% indicates values of less than 1%.

**Seasonal Prevalence**

The seasonal prevalence of all parasites was highest (43-51%) between December and March but lower during the rest of the year reaching a low of 34 - 36% in August and September. This pattern was clearly influenced by a corresponding pattern in the seasonal prevalence of *B. hominis*, the most common parasite, which reached 20 - 23% then declined to 13 - 15%, during the same time periods, respectively (Table 3). The seasonal prevalence of *Cryptosporidium parvum* infections was more or less stable throughout the year (4 - 7%) but declined to a low of 3% in August. The prevalence of *D. fragilis* was highest in February (3%) and November (4%). The prevalence of *E. coli* cysts and trophozoites was highest in July (8%) but lower (3 - 4%) between December and May. The prevalence of each of *E. Hartmanni* and *E. histolytica/dispar* cysts and trophozoites was highest (10%) in February and lowest (2%) in November. The prevalence of the amoeba trophozoites was usually relatively higher than that of cysts during most months. The numbers of detected *G. intestinalis* were too small to produce credible seasonal data (Table 3).

Parasite species	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	TOTAL
Blastocystis hominis	75(23)*	72(21)	73(22)	80(20)	63(16)	62(18)	53(18)	41(13)	47(15)	58(19)	58(20)	50(20)	732(19)
Cryptosporidium parvum	17(5)	18(5)	20(6)	16(4)	22(6)	19(6)	12(4)	9(3)	22(7)	17(6)	13(5)	16(6)	201(5)
Dientamoeba fragilis	4(1)	11(3)	4(1)	9(2)	5(1)	4(1)	1(0)	2(1)	1(0)	7(2)	11(4)	1(0)	60(2)
Entamoeba coli C**	6(2)	8(2)	3(1)	8(2)	8(2)	9(3)	12(4)	6(2)	5(2)	6(2)	8(3)	6(2)	85(2)
E. coli T**	7(2)	3(1)	5(2)	7(2)	3(1)	6(2)	12(4)	5(2)	2(1)	8(3)	10(3)	3(1)	71(2)
E. hartmanni C	11(3)	13(4)	11(3)	8(2)	10(3)	10(3)	7(2)	11(4)	5(2)	5(2)	3(1)	9(4)	103(3)
E. hartmanni T	10(3)	20(6)	9(3)	12(3)	9(2)	10(3)	9(3)	10(3)	12(4)	8(3)	4(1)	10(4)	123(3)
E. histolytica/ dispar C	6(2)	10(3)	8(2)	2(1)	10(3)	10(3)	8(3)	4(1)	4(1)	5(2)	2(1)	6(2)	75(2)
E. histolytica/ dispar T	15(5)	24(7)	10(3)	14(4)	15(4)	19(6)	15(5)	19(6)	10(3)	6(2)	3(1)	9(4)	159(4)
Giardia intestinalis C	2(1)	1(0)	1(0)	3(1)	2(1)	0(0)	1(0)***	2(1)	3(1)	2(1)	0(0)	1(0)	18(0)
G. intestinalis T	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	1(0)	0(0)	1(0)	0(0)	2(0)
Total # of infections	153	180	0	0	0	0	0	0	0	0	0	0	0
Total examined	322	348	333	397	386	339	301	312	307	301	286	251	3883
Prevalence	47%	51%	43%	40%	38%	43%	43%	34%	36%	40%	39%	44%	41%

**Table 3:** Seasonal distribution of infections with major protozoan parasites diagnosed from 3883 patients examined from Los Angeles County between 1996 and 2013.

\*No. of infections (% of examined patients per each month). Percentages are rounded.

\*\*C=cysts, T=-trophozoites.

\*\*\* 0% indicates values of less than 1%.

**Discussion**

**Prevalence in the USA**

The overall prevalence of infection with all investigated protozoans was 41% of double fecal samples examined between 1996 and 2013 from 3,883 patients in the Los Angeles County. Infections with *B. hominis* made up roughly half (45%) of all protozoans studied and

noted in Table 1. Los Angeles is an urban/suburban area and a 41% prevalence rate is markedly higher than the 32% and 36% prevalence rates reported earlier in the United States and California, respectively [2]. The testing procedures employed in this study produced a prevalence rate of 32.6% (3,373 infected of 10,358 examined patients throughout the United States) between 1996 and 1998 [8]. An almost identical prevalence of 32% was reported in our comprehensive study of the prevalence of intestinal parasites in 5,792 fecal specimens from 2,896 patients in the United States [2]. Our results reflect our most efficient methods of parasite detection [8] which show a considerably higher prevalence rates than others across the country. For instance, Kappus [4] reported US prevalence of 20% (from 216,275 stool specimens) compared with 19.7% (from 178,786 stool specimens) reported by state diagnostic laboratories in 1987. Similarly, Garcia and Brukner [12] reported a prevalence of 20.6% from 2,360 US patients. Differences in test populations or in the composition of the component parasite may be involved. Church., *et al.* [5] reported a low prevalence of 6.4-7.2% of 2,604 fecal specimens from Colorado, Montana, New Mexico, and Utah were infected with parasites between August, 2006 and April, 2007. Amin [2] reported 19%, 50%, 39%, and 29% from the same states in the same order. Church., *et al.* [5] attributed their low prevalence rates, in part, to their inability to detect infections with *C. parvum* and *C. cayetanensis*; Quest Diagnostics, Denver tested their specimens.

### Prevalence in Developing Countries

In comparable urban/suburban areas in Africa, Asia and South America, the prevalence of parasitic infections was mostly similar to ours (Table 1) but occasionally somewhat lower or markedly higher. It was 21.4% of 5,990 patients in Madhya Pradesh, India [13], 23.14% of 350 patients in Dhakka University, Bangladesh [14], 29.26% of 287 patients in Muzaffaraband, Pakistan (Chaudhry., *et al.* [15], who reviewed prevalences in 14 Pakistani cities from 40,096 subjects), 33% of 199 patients in Chennai, India [16], 33.4% of 1,127 patients in Izmir, Turkey [17], 42.9% of 2,400 patients in Kumasi, Ghana [18], 47% of 293 patients in Varamin, Iran [19], 47.2% of 1267 patients in León, Nicaragua [20], 50.5% of 93 patients in Central Nigeria [21], 62% of 195 patients (protozoans only) in Tamil Nadu, India [22], and 75.1% of 1,227 patients in Bioko, Equatorial Guinea [23]. On rare occasions, the overall prevalence of intestinal infection was very low reaching 5.92% of 5,743 patients in Eghbalieh City, Iran [24]. We believe such differences in prevalence to be attributable to demography, diet, environmental exposure, social habits, urbanization, and zoonotic relationships as can be discerned from the above articles.

### Prevalence in Developed Countries

Fletcher., *et al.* [12] summarized 33 prevalence studies from cities in developed countries, 23 in Europe and 10 from the US, Canada, Australia, and Korea. The prevalence rate of *B. hominis* varied between 1% in Danish counties, Denmark to 16.9% in Sydney, Australia. For *C. parvum*: between 0.4% in Melbourne, Australia to 9.1% in Helsinki, Finland. For *E. histolytica/dispar*: between 0.4% in Helsinki to 3.5% in Noumea, New Caledonia. For *G. intestinalis*: between 0.3% in Melbourne, Australia to 29% in Helsinki. *Entamoeba coli* was reported only once from Brussels, Belgium at a rate of 5.4% and *D. fragilis* 6 times between 0.4% in Melbourne, Australia and 14.6% in Holland.

### Prevalence of Individual Protozoan Species

In the United States, *B. hominis* was the most dominant protozoan parasite. Its reported prevalence was 19% (45% of all infections) (Table 1), 23% in 2000 [2], 20 - 30% [25], and 12.2% [12]. It was also the most dominant parasite species reported by Kappus [4] and Church., *et al.* [5] but at surprisingly lower prevalence of only 2.6% and 4.3%, respectively. In developed countries, *B. hominis* appears to also be the dominant intestinal parasite, e.g. Izmir, Turkey [17], Amsterdam, Holland [26] (24.2%), Sydney, Australia [27] (18 - 21%), Thessaloniki, Greece [28] (5.3 - 16.8%), Stockholm, Sweden [29] (4.0%), Berlin, Germany [30] (7.6%), Helsinki, Finland [11] (Table 1; 13%), Rome, Italy [31] (7.5 - 14.1%), and Brussels, Belgium [11] (Table 1; 9.8%). In most developing countries in Asia, Africa and South America, however, *E. histolytica/dispar* and /or *G. intestinalis* appear to be the dominant parasites, e.g., Jordan [32]; India [11,13,33]; Saudi Arabia [34]; Lebanon [35], Nicaragua [20], Ghana [18], Equatorial Guinea [23], Pakistan [15], Iran [19], and Bangladesh [14]. Fletcher., *et al.* [11] concluded that "while some enteric protozoa, such as *Entamoeba* sp., *Cryptosporidium*, and *Giardia* are isolated frequently from diarrheal patients in developing regions such as Asia and sub-Saharan Africa, others, such as *Blastocystis* spp. and *Dientamoeba fragilis* are isolated mainly in developed countries" We concur.



In the present study, the next highest prevalence to *B. hominis* were 6%, 5%, 5%, and 41% noted for *E. histolytica/dispar*, *E. hartmanni*, *C. parvum*, and *E. coli.*, respectively. Only 20 patients (< 1%) were infected with *G. intestinalis* (Table 1). This ranking was not consistent in some other studies in the United States. For example, in the Rocky Mountain states, the prevalence of *Endolimax nana* and *G. intestinalis* ranked second (1.5%) and third (1.4%) to *Blastocystis* infections [5]. The prevalence of *E. histolytica/dispar* of 6% (Table 1) is markedly higher than the 0.9% reported in a large 1987 survey than the estimated prevalence of 4% in the United States [36]. The prevalence of *C. parvum* of 5% (Table 1) is higher than 0.6 - 4.3% reported elsewhere in North America but less than the 3 - 20% known from other area of the world (Asia, Australia, Africa, and Central and South America) [10]. *Cryptosporidium parvum* appears to be underdiagnosed in the western hemisphere; its seroprevalence in Europe and North America is usually between 25% and 35% and may reach 64% in South America [37]. *Cryptosporidium* oocysts were observed in 27% of drinking water sampled from 66 surface water treatment plants in 14 states and one Canadian province [38]. Differences in the prevalence and composition of the intestinal parasite fauna in different geographies are probably attributable to demography, diet, environmental exposure, social habits, urbanization, and zoonotic relationships as well as to the structure of the edaphic conditions and weather affecting the extra-human stages of the parasites particular to each location.

### Annual Prevalence

The total number of samples submitted from the Los Angeles area was highest in 1996 - 1997 but declined then stabilized afterwards corresponding with changing patterns of patient traffic. The prevalence of most parasitic infections was highest during 1996 - 97 (63%) then gradually declined reaching 21% in 2012 - 2013 agreeing with that of *B. hominis*, the most common parasite detected, being 21% and 7%, in the same order (Table 2). The prevalence of all other protozoans, except *C. parvum*, was highest in 1996 - 1997. The prevalence of *C. parvum* progressively increased from 2% in 1996-1997, when prevalence of all other infections were at a minimum, to a high of 11% in 2008-2009 then declined afterwards. *Giardia intestinalis* was the only protozoan that was consistently identified in 1% or less of the samples throughout the study period.

### Decline in Annual Prevalence

The general decline in the prevalence of all parasites and especially of the dominant *B. hominis* over the years was similar to declines over time reported in other studies. In 10,582 fecal specimens from U.S. general population [3], reported declining *B. hominis* prevalence rates of 23%, 20%, 15%, and 11% between 2000 and 2004, respectively. A similar study of *C. parvum* from 9,856 fecal specimens from US general population between 2003 and 2005 noted an almost even prevalence of 5 - 6% [39]. Annual prevalence rates of microsporidiosis from fecal specimens of 8,550 HIV-infected patients in Southern California demonstrated a decline from 8.8% in 1993, 9.7% 1994, 6.6% in 1995, and 2.9% in 1996 which was attributed to "the use of multi-drug antiretroviral regimens and the use of protease inhibitors, a new class of antiretroviral agents, the first of which was licensed in 1995" [6]. In Izmir, a Turkish Mediterranean coastal city with a climate similar to that of Los Angeles, the prevalence of intestinal parasites was 42.5% in 2003 [7] and 65% five years earlier in the same area [40]. Of 18,563 hospital patient records studied in Qatar, the prevalence of intestinal parasites decreased from 13.4% in 2005 - 2008 to 6.6% in 2009-2011 [41]. During this period, the prevalence of *B. hominis*, *G. intestinalis*, and pathogenic amoeba decreased from 4.3% to 2.9%, 1.9% to 1.4%, and 0.29% to 0.25%, respectively. This decline was attributed, in part, to improved screening of foreign workers. In Madhya, Pradesh, India, the prevalence of intestinal parasites (mostly *Giardia* and *E. histolytica/dispar*), during 2003, 2006, 2007, 2008, 2009, 2010, and 2011 was 59.5% [42], 24.1%, 22.3%, 20.3%, 19.9%, and 20.4%, 21.4% [13], respectively. In Lebanon, Araj., *et al.* [35] reported decreasing prevalence from 14% to 12% in *E. histolytica/dispar* and from 16% to 6% in *G. intestinalis* between 1997-1998 and 2007-2008 in 14,771 and 7,477 fecal specimens tested, respectively. We can attribute the overall decline in prevalence of parasitic protozoan infections over time to improved health education, better preventive measures, and more effective drug therapies.

### Seasonal Prevalence

The seasonal prevalence of all infections in our study populations did not show a dramatic seasonal periodicity [2,3,39]. However, in the present investigation, the seasonal prevalence of all protozoan infections was highest in February (51%) and lowest in August (34%) corresponding with the pattern in *B. hominis* (Table 3). In 2000, we found that the seasonal prevalence of all infections from 48 states and

the District of Columbia was highest in September and October (42% and 43%) and lowest in February (22%) [2]. In our studies of 2000 [2], and 2002-2004 [3], the seasonal prevalence of *B. hominis* in the general US populations also did not show any marked seasonality but was highest in September (23%) and lowest in February (13%). The seasonal prevalence of *C. parvum* in the Los Angeles area (this paper) was more or less stable throughout the year (3% in August to 7% in September). The prevalence of microsporidiosis in 8,439 fecal specimens from persons with diarrhea and human immunodeficiency syndrome in southern California showed no seasonal variation [6]. The prevalence of *C. parvum* from throughout the US was low (3 - 6%) during the colder months of the year (October to March) but higher during the warmer months of April and May, reaching 9% of 9,856 fecal specimens examined between 2003-2005 [39]. "Peaks in *C. parvum* prevalence appear to correspond with warmer seasons in temperate and tropical climates especially when associated with rain. During the rainy seasons, the run off from cattle farms readily contaminate surface waters feeding into water treatment plants as happened during the March-April, 1993 Milwaukee outbreak [39]. This pattern of seasonal waterborne fecal contamination has also been reported by other observers throughout the world, e.g. US [43], New Orleans [44], Peru [45], England [46], Korea [47], Uganda [48], Jordan [49], Guatemala [50], Indonesia [51], and Zambia [52]. *Cryptosporidium* infections from 2000 HIV-positive patients in Benin City, Nigeria were reported to also be associated with the rainy season [53]. The infectious stages of these soil-based intestinal parasites released into the environment are clearly vulnerable to seasonal variations in temperature, rainfall and humidity before they encounter other hosts. Transmission depends on the production of and host encounters with parasite stages in the environment. Seasonal variations in host immune system being weaker in the winter may also be involved [54].

### Seasonality of Soil-Based Infections

The seasonal prevalence of other soil-based infections, e.g., *E. histolytica/dispar* and *G. intestinalis* was also highest in the rainy season and lowest in winter or summer in 350 fecal specimens tested in Dhakka University Medical Center, Bangladesh [14]. Similar results were reported for the same two protozoans from 23,278 fecal specimens in Qassim region of Saudi Arabia and from 22,970 stools in the Nablus area, Jordan by Imam, *et al.* [34] and Ali-Shtayeh, *et al.* [32], respectively. Patel [33], however, reported "no significant" seasonal differences in the prevalence of *E. histolytica/dispar*, *G. intestinalis*, and *E. coli* from 36,000 reports of inpatients and outpatients of Bombay hospital, India analyzed between 1966 and 1975.

Non-water sources of seasonal *Cryptosporidium* infections (food stuffs, drinks, animal to person, person to person contact, exposure to contaminated recreational water, among others) are known in arid desert countries mostly in the coldest season and may reach as high as 50% of total exposures [55]. This pattern is best illustrated in the desert country of Kuwait where 77% of the infections occur between November and April [56]. Seasonal prevalence of other protozoans investigated (Table 3) were more or less stable throughout the year.

\*This article is based on the November, 2014 proceedings of presentations made at the PCI-Europe workshops at the Royal Society of Medicine Conference Center, the Bowskill Clinic, and the Kings College in London, UK on March 5-8, 2014.

### Bibliography

1. Amin OM. "Prevalence and host relationships of intestinal protozoan infections during the summer of 1996". *Explore* 8.2 (1997): 29-35.
2. Amin OM. "Seasonal prevalence of intestinal parasites in the United States during 2000". *American Journal of Tropical Medicine and Hygiene* 66.6 (2002): 799-803.
3. Amin OM. "The epidemiology of Blastocystis hominis in the United States". *Research Journal of Parasitology* 1.1 (2006): 1-10.
4. Kappus KD, *et al.* "Intestinal parasitism in the United States: update on a continuing problem". *American Journal of Tropical Medicine and Hygiene* 50.6 (1994): 705-713.

5. Church C., *et al.* "Intestinal infections in humans in the Rocky Mountain region, United States". *Journal of Parasitology* 96.1 (2010): 194-196.
6. Contreas CN., *et al.* "Examination of the prevalence and seasonal variation of intestinal microsporidiosis in the stools of persons with chronic diarrhea and human immunodeficiency virus infections". *American Journal of Tropical Medicine and Hygiene* 58.5 (1998): 559-561.
7. Senay H and Macpherson D. "Blastocystis hominis epidemiology and natural history". *Journal of Infectious Diseases* 162.4 (1990): 987-990.
8. Amin OM. "Evaluation of a new system for the fixation, concentration, and staining of intestinal parasites in fecal specimens, with critical observations on the trichrome stain". *Journal of Microbiological Methods* 39.2 (2000): 127-132.
9. Tan KSW. "New insights on classification, identification, and clinical relevance of Blastocystis spp". *Clinical Microbiology Reviews* 21.4 (2008): 639-666.
10. Garcia LS. "Diagnostic medical Parasitology, 2001". American Society Microbiology Press, Wash, DC. (2001): 1092.
11. Fletcher SM., *et al.* "Enteric Protozoa in the developed world". *Clinical Microbiology Reviews* 25.3 (2012): 420-449.
12. Garcia LS., *et al.* "Clinical relevance of Blastocystis hominis (letter)". *Lancet* 323.8388 (1984): 1233-1234.
13. Marothi Y and Singh B. "Prevalence of intestinal parasites at Ujjain, Madhya Pradesh, India: Five-year study". *African Journal of Microbiology Research* 5.18 (2011): 2711-2714.
14. Khanum H., *et al.* "Occurrence of intestinal parasites among the teachers, students and staffs of Dhaka University". *Journal of the Asiatic Society of Bangladesh, Science* 39.2 (2013): 239-246.
15. Chaudhry ZH., *et al.* "Epidemiological factors affecting prevalence of intestinal parasites in Children of Muzaffarabad District". *Pakistan Journal of Zoology* 36.4 (2004): 267-271.
16. Fernandez MC., *et al.* "A comparative study of the intestinal parasites prevalent among children living in urban and rural settings in and around Chennai". *Journal of Communication Disorders* 34.1 (2002): 35-39.
17. Aksoy Ü., *et al.* "Demographic status and prevalence of intestinal parasitic infections in schoolchildren in Izmir, Turkey". *Turkish Journal of Pediatrics* 49.3 (2007): 278-282.
18. Walana W., *et al.* "Prevalence of intestinal protozoan infestation among primary school children in urban and peri-urban communities in Kumasi, Ghana". *Science Journal of Public Health* 2.2 (2014): 52-57.
19. Aminzadeh Z., *et al.* "Prevalence of intestinal parasites and related factors in primary school children in Varamin". *Iranian Journal of Pediatrics Society* 1.2 (2007): 55-58.
20. Téllez A., *et al.* "Prevalence of intestinal parasites in the human population of León, Nicaragua". *Acta Tropica* 66.3 (1997): 119-125.
21. Ikeh E., *et al.* "Intestinal parasitism in rural and urban areas of North Central Nigeria: An update". *Internet Journal of Microbiology* 2.1 (2005): 1-11.



22. Rayan P, *et al.* "Geographical location and age affects the incidence of parasitic infestations in school children". *Indian Journal of Pathology and Microbiology* 53.3 (2010): 498-502.
23. Roche J and Benito A. "Prevalence of intestinal parasite infections with special reference to *Entamoeba histolytica* on the Island of Bioko (Equatorial Guinea)". *American Journal of Tropical Medicine and Hygiene* 60.2 (1999): 257-262.
24. Sadeghi H, *et al.* "Prevalence of intestinal parasites in a population in Eghbalieh City from Qazvin Province, Iran". *Journal of Parasitic Diseases* 39.2 (2013): 126-129.
25. Lee MJ. "Parasites, yeasts and bacteria in health and disease". *Journal of Advances in Medicine* 8 (1995): 121-130.
26. Bart A, *et al.* "Diagnosis and subtype analysis of *Blastocystis* sp. in 442b patients in a hospital setting in the Netherlands". *BMC Infectious Diseases* 13 (2013): 389.
27. Stark D, *et al.* "Prevalence of enteric protozoa in human immunodeficiency virus (HIV) positive and HIV negative men who have sex with men from Sydney, Australia". *American Journal of Tropical Medicine and Hygiene* 76.3 (2007): 549-552.
28. Papazahariadou MG, *et al.* "Prevalence of gastrointestinal parasites in the Greek population: local people and refugees". *Annals of Gastroenterology* 17.2 (2004): 194-198.
29. Svenungsson B, *et al.* "Enteropathogens in adult patients with diarrhea and healthy control subjects: a 1-year prospective study in a Swedish clinic for infectious diseases". *Clinical Infectious Diseases* 30.5 (2000): 770-778.
30. Jansen A, *et al.* "Aetiology of community-acquired, acute gastroenteritis in hospitalized adults: a prospective cohort study". *BMC Infectious Diseases* 8 (2008): 143.
31. Masucci L, *et al.* "Intestinal parasites isolated in a large teaching hospital, Italy, 1 May 2006 to 31 December, 2008". *Euro Surveillance* 16.24 (2011): 19891.
32. Ali-Shtayeh MS, *et al.* "Prevalence and seasonal fluctuations of intestinal parasitic infections in the Nablus area, West Bank of Jordan". *Annals of Tropical Medicine and Parasitology* 83.1 (1989): 67-72.
33. Patel JC. "Ten year study of stool samples with particular reference to intestinal parasites". *Journal of Postgraduate Medicine* 32.4 (1986): 219-224.
34. Imam A, *et al.* "Frequency and seasonality of intestinal parasitism in Qassim region, Saudi Arabia". *Pakistan Journal of Medical Sciences* 28.5 (2012): 913-916.
35. Araj GF, *et al.* "Trends and prevalence of intestinal parasites at a tertiary care center in Lebanon over a decade". *Le Journal Medical Libanais* 59.3 (2011): 143-148.
36. Kelsall BL and Ravdin JI. "Amebiasis: Human infection with *Entamoeba histolytica*". Tsieh S, ed. *Progress in Clinical Parasitology*. Ann Arbor, MI: CRC Press. 4 (1994): 27-54.
37. Current WL and Garcia LS. "Cryptosporidiosis". *Clinics in Laboratory Medicine* 11 (1991): 873-895.
38. LeChevallier MW, *et al.* "Giardia and *Cryptosporidium* spp. in filtered drinking water supplies". *Applied and Environmental Microbiology* 57.9 (1991): 2617-2621.

39. Amin OM. "Epidemiology of *Cryptosporidium parvum* in the United States". *Parasitology Unit Journal* 1 (2008): 15-22.
40. Akisu C., *et al.* "Investigation of intestinal parasites in school children living under low socio-economic conditions in Izmir". *Acta Parasitology* 24 (2000): 52-54.
41. Abu-Madi MA., *et al.* "Intestinal parasitic infections among long-term-residents and settled immigrants in Qatar in the period 2005 to 2011". *American Journal of Tropical Medicine and Hygiene* 88.6 (2013): 1185-1195.
42. Rao VG., *et al.* "Intestinal parasitic infections, anaemia and undernutrition among tribal adolescents of Madhya Pradesh". *Indian Journal of Community Medicine* 27 (2003): 26-29.
43. Hlavsa MC., *et al.* "Cryptosporidiosis surveillance- United States 1999-2002". *Morbidity and Mortality Weekly Report. Surveillance Summaries* 54.1 (2005): 1-8.
44. Inungu JN., *et al.* "Risk factors, seasonality, and trends of cryptosporidiosis among patients infected with human immunodeficiency virus". *American Journal of Tropical Medicine and Hygiene* 62.3 (2000): 384-387.
45. Bern C., *et al.* "Epidemiologic differences between cyclo-sporiasis and cryptosporiasis in Peruvian children". *Emerging Infectious Diseases* 8.6 (2002): 581-585.
46. Howe AD., *et al.* "Cryptosporidium oocysts in a water supply associated with a cryptosporidiosis outbreak". *Emerging Infectious Disease* 8.6 (2002): 619-624.
47. Chai JY., *et al.* "High prevalence and seasonality of cryptosporidiosis in a small rural village occupied predominantly by aged people in the Republic of Korea". *American Journal of Tropical Medicine and Hygiene* 65.5 (2001): 518-522.
48. Tumwine JK., *et al.* "Cryptosporidium parvum in children with diarrhea in Mulago hospital, Kampala, Uganda". *American Journal of Tropical Medicine and Hygiene* 68.6 (2003): 710-715.
49. Mahgoub ES., *et al.* "Cryptosporidiosis in children in a north Jordanian paediatric hospital". *Eastern Mediterranean Health Journal* 10 (2004): 494-501.
50. Bern C., *et al.* "The contrasting epidemiology of Cyclospora and Cryptosporidium among outpatients in Guatemala". *American Journal of Tropical Medicine and Hygiene* 63.5-6 (2000): 231-235.
51. Katsumata T., *et al.* "Cryptosporidiosis in Indonesia: a hospital based study and a community- based survey". *American Journal of Tropical Medicine and Hygiene* 59.4 (1998): 628-632.
52. Nchito M., *et al.* "Cryptosporidiosis in urban Zambian children: an analysis of risk factors". *American Journal of Tropical Medicine and Hygiene* 59.3 (1998): 435-437.
53. Akinbo FO., *et al.* "Seasonal variation of intestinal parasitic infections among HIV-positive patients in Benin City, Nigeria". *Ethiopian Journal of Health Sciences* 21.3 (2011): 191-194.
54. Altizer S., *et al.* "Seasonality and dynamics of infectious diseases". *Ecology Letters* 9.4 (2006): 467-484.
55. Anonymous. "Cryptosporidium and water- the numbers game". *Hlth Stream*. 16 (1999): 1- 4.

56. Gaitai W, *et al.* "Cryptosporidiosis: prevalence, genotype analysis, and symptoms associated with infections in children in Kenya". *American Journal of Tropical Medicine and Hygiene* 75.1 (2006): 78-82.

**Volume 2 Issue 6 May 2017**

**© All rights reserved by Omar M Amin and Karim O Amin.**