

Status of Skilled Birth Attendance Utilization and Determinants Among Women of Child Bearing Age in Chencha Woreda, Gamo Gofa Zone, Southern Ethiopia, December 2016

Mende Mensa^{1*} and Melkenesh Nigatu Belijo²

¹Lecturer and Researcher, Arba Minch College of Health Sciences, Arba Minch, Ethiopia

²Arba Minch General Hospital, Arba Minch, Ethiopia

*Corresponding Author: Mende Mensa, Lecturer and Researcher, Arba Minch College of Health Sciences, Arba Minch, Ethiopia.

Received: September 02, 2017; Published: September 27, 2017

Abstract

Background: Approximately 99% of maternal deaths are occurring in developing countries and most of these deaths could have been prevented if mothers had access antenatal care (ANC) during pregnancy, were attended by a skilled birth attendant. However the prevalence of skilled birth attendance and institutional delivery were still low in developing countries like Ethiopia. Therefore this study was aimed at exploring the reasons for low skilled birth attendance in rural communities of Chencha District Gamo Gofa Zone, Southern Ethiopia.

Objective: To assess the status of skilled birth attendance and its determinants among mothers in rural communities of Chencha district, southern Ethiopia, December 2016.

Methods: A community based cross sectional study that employed quantitative data collection method was conducted. Multi stage sampling technique was used for selection of study participants. A pre tested semi-structured questionnaires via interview was used to collect data on different variables.

Results: There are 500 women who had children less than 5 years old were participated in this study. Majority of participants 356 (71.2%) were in the age group of 26 - 30 years with Mean age of $27.44 \pm (3.076)$ years. Age of the mother 21 - 25 years [AOR = 1.56 (1.204 - 3.516), Government employed husband [AOR = 1.669 (1.475 - 81.642), Mothers with monthly income above 5000 ETB [AOR = 2.196 (1.051 - 4.590), No ANC follow-up [AOR = 0.915 (0.375 - 0.953), Mothers who reported health professionals are not well skilled [AOR = 0.227 (0.085 - 0.607) and Health facility in less than 5 km distance [AOR = 3.116 (1.916 - 5.068) were independent predictors of skilled birth attendance utilization.

Conclusions: Our study revealed that prevalence of skilled birth attendance below one third. Age of the mother 21 - 25 years, Mothers with government employed husband, mothers with monthly income above 5000 ETB, ANC follow-up, Health professionals skilled and Health facility in less than 5km distance were predictors of skilled birth attendance utilization. Designing strategies to address these variables by responsible bodies is critical.

Keywords: Skilled Birth Attendance; Knowledge; Attitude; Antenatal Care; Southern; Ethiopia

Abbreviations

ANC: Antenatal Care; BEmONC: Basic Emergency Obstetrics and Neonatal Care; C/S: Cesarean Section; CEmON: Comprehensive and Emergency Obstetrics and Neonatal care; CI: Confidence Interval; DHS: Demographic and Health Survey; EDHS: Ethiopian Demographic and Health Survey; EmOC: Emergency Obstetric Care; ETB: Ethiopian Birr; HEW: Health Extension Worker; HSDP: Health Sector Development Program; MDG: Millennium Development Goal; MMR: Maternal Mortality Ratio; MOH: Ministry of Health; PAD: Professionally Assisted Delivery; PI: Principal Investigator; RH: Reproductive Health; SBA: Skilled Birth Attendants; SNNPR: Southern Nation, Nationalities and Peoples Region; SPSS: Statistical Package for Social Studies; SSA: Sub Sahara Africa; SVD: Spontaneous Vaginal Delivery; WCBA: Women of Child Bearing Age; WHO: World Health Organization

Introduction

Maternal health encompasses social, cultural, health systems and health policy factors. Among these factors complications of pregnancy and childbirth are a leading cause of maternal morbidities and mortalities for women of reproductive age (15 - 49 years) in developing countries [1,2].

Nearly 3.4 million of the 8 million infant deaths each year occur within the first week of life and are often due to a lack of or inappropriate care during pregnancy, delivery and the post-partum period [3]. Hemorrhage, infection, eclampsia, obstructed labour and post abortion complications accounts for 80% of all direct causes of maternal deaths, and the final 20% are due to indirect obstetric causes like HIV/AIDS, anaemia, malaria and cardiovascular disease [4].

Approximately 303,000 maternal deaths occurred globally in 2015, yielding an overall MMR of 216 maternal deaths per 100,000 live births. The global lifetime risk of maternal mortality is approximately 1 in 180 for 2015. The overall MMR in developing regions is 239, which is roughly 20 times higher than that of developed regions. Developing regions account for approximately 99% (302,000) of the estimated global maternal deaths, with sub-Saharan Africa alone accounting for roughly 66% (201,000). The lifetime risk of maternal mortality is estimated at 1 in 36 in sub-Saharan Africa, contrasting sharply with approximately 1 in 4900 in developed countries. In Ethiopia in 2016 the level of maternal mortality (MMR) is high 412 per 100,000 live births [5,6].

Global targets for ending preventable maternal mortality (EPMM) By 2030, every country should reduce its maternal mortality ratio by at least two thirds from the 2010 baseline, and no country should have an MMR higher than 140 deaths per 100 000 live births (twice the global target) [7].

In low and middle income countries, a significant proportion of women still deliver at home and more than 80% of maternal deaths worldwide are due to five direct causes, hemorrhage, sepsis, unsafe abortion, obstructed labor, and hypertensive diseases. Despite the fact attending antenatal care during pregnancy and giving birth in health facility could prevent most of maternal deaths [8-10].

A recent study conducted in Benin, Ecuador, Jamaica and Rwanda demonstrated that higher levels of skilled attendants during delivery lead to lower maternal mortality. It is estimated that, if there were SBAs at all deliveries, maternal mortality could be reduced by 13 - 33%. Utilization of maternal health services in developing countries is low and hindered by factors like lack of infrastructure, quality of services provided and accessibility of the service [11-13].

Study conducted in North West Ethiopia showed that only 38.2% of women gave birth in health facilities for their most recent birth while the most 223 (61.8%) delivered at home. The major reasons of institutional delivery were fear of complications (77.5%), to get better service (56.5%) safe and clean delivery (46.4%) and they were communicated to deliver at health facilities during ANC attendance, 47(35.5%) [14].

Mothers' educational status, family size and household wealth are highly correlated with whether their deliveries are assisted by a skilled provider and whether they are delivered in a health facility [15-17].

Studies conducted in Ethiopia indicated that mothers with lower age at first pregnancy, having more than one previous pregnancy, with in more than one birth order, received ANC service, birth complications in previous and immediate pregnancies and had every used professionally assisted delivery in preceding delivery were commonly found to be significantly associated with safe delivery service utilization [18,19].

Similar Studies showed that Mothers with the age range of 15 - 19 years, with parity 1, married to husbands those completed secondary school, who had received four antenatal care, mothers, who had ever given birth at least once earlier in health facility and those having known risk of giving birth at home were more likely to utilize skilled birth attendances than their counter parts [20-22].

Regardless of the government's efforts to expand health service facilities and promote institutional delivery service in Ethiopia, most of the mother gives birth at home. The prevalence of live births delivered by a skilled provider was 28% and the prevalence of institutional delivery was 26% in the 2016 EDHS. This is much lower in the rural communities 21% of births were assisted by a skilled provider and 20% were delivered in a health facility [23-27].

In Ethiopia, studies addressing determinants of maternity care services utilization were limited particularly in Gamo Gofa zone. Additionally, little is known about the determinants of utilization of skilled birth attendances services in the study area. Therefore this study was designed to fill these discrepancies by assessing the determinants of utilization of skilled birth attendances among women in child-bearing age at rural villages in Chencha district Gamo Gofa Zone, Southern Ethiopia. The study findings will have input on maternal health improvement in the region as well as in the study area. The findings will also be used by responsible bodies who work on reproductive health services to address issues of reducing maternal morbidity and mortality.

Methods and Materials

Study Area and period

The study was conducted in Chencha district from December 01 to 30, 2016. The district is one of the 15 districts in Gamo Gofa zone, Southern Nations and Nationalities Regional State. Based on projection from the 2007 E.C population and housing Census the woreda is estimated to be 142,063. The district has 48 kebeles; 3 town administration namely Chencha town, Ezzo town and dorze town which constitute about 14% of the woreda population and the remaining 86% of the population resides in the rural community.

Study design: A community based quantitative cross-sectional study design was employed.

Source and Study population

Source population: All women in childbearing age who were permanent residents at least for one year in the study area were source population.

Study population: All women in childbearing age that had given birth in the five years before the survey residing in randomly selected kebeles.

Study subjects: Women of childbearing age those had given birth in the past five years preceding the study period proportionally allocated in selected kebeles.

Sample size determination and Sampling technique/sampling procedure

Sample size determination and Sampling technique

In the survey, multistage sampling technique was used. The following assumption and formula were considered in calculating the size of the sample. Proportion (p = 26.8%) from study conducted in Sidama Zone" [22], at 95% confidence interval, 10% non-response rate and a design effect of 1.5 were considered. Finally, 500 sample women of childbearing age were drawn. The sample was calculated using single Population proportion formula.

$$N = \frac{Z_{\alpha/2}^2 * P(1 - P)}{D^2} = 333$$

Where:

N = sample size of women who gave birth at health institution by skilled attendants

Z α /2 = is the corresponding Z value to 95% significance level = 1.96

P = Proportion of women who had given birth at health facilities by skilled attendants = 26.8%.

d = degree of freedom = 5%.

The Final sample size = 498.2 \approx 500

Sampling procedure

Among forty five rural kebeles in the woreda five rural kebeles were selected randomly. From the selected kebeles, households with women who had given birth in last five years preceding the survey were identified by using the registration at health post. The total sample size had been then proportionally allocated from total women of childbearing age in each kebele. In cases where a woman had given birth more than once, the most recent pregnancy was considered for the present study. Based on this, a sampling frame that enlists all eligible mothers had been prepared and 500 women were included in the study using probability sampling technique.

Eligibility criteria

Inclusion criteria: Reproductive age women who are residing in the study area for at least one year and have given birth within five years preceding the survey.

Exclusion criteria: Women who were mentally and physically non-capable and unwilling to participate.

Variables of the study

Dependent Variable: Utilization of skilled birth attendance (SBA) services.

Independent Variables

Socio demographic variables: Maternal age, marital status, age at marriage, ethnicity, religion, income, Family size, residence, educational status, occupational status.

Obstetrics characteristics: Age at first pregnancy, birth order, frequency and use of antenatal care, child birth complications, Women's knowledge and attitude.

Health facility factors: Distance of health facility from their home, availability of health facility and health care providers.

Data collection procedure

Face-to-face interview with semi-structured questionnaire that was adapted from Ethiopian demographic and health survey and related thesis works [16]. Women in the age range of childbearing had been interviewed by going house to house in the randomly selected kebeles. The English version of the questionnaire was translated into a local language, Gamogna, by experts and translated back into English for consistency. Three diploma nurses were recruited and oriented for one day by the principal investigator prior to data collection. One data collector was assigned per kebele. One supervisor with a health background holding a first degree was assigned per kebele in order to supervise and assist the data collection process. Mothers who were not present during the first visit had been revisited twice and the result of the visit was recorded on the tool.

Data quality management

The questionnaire was pre-tested on 5% of the respondents in Doko shaye village that was randomly selected for this purpose. The kebele is similar with socio-demographic characteristics with the people in the study areas. Findings had been discussed among data collectors and supervisors and the tool was amended before actual data collection based on findings of the pre-test. Data quality had been ensured during collection, coding, entry and analysis. During data collection, adequate orientation and follow-up had been provided to data collectors and supervisors. Supervision of data collectors included observation of how the data collectors were going to administer questions. Supervisors checked the filled questionnaires on a daily basis for completeness, clarity. Then, the principal investigator had double-checked randomly the questionnaire for the completion each day.

Data processing and analysis

Finally, the data had first been checked manually for completeness then recoded, edited and entered into Epi data version 3.1 and exported to SPSS version 20.0 for further analysis. The data had further been cleaned by visualizing, calculating frequencies and sorting. After exploration, bivariate and multivariate analyses had been computed. The bivariate analysis using logistic regression was done to see the crude association between dependent and independent variables so that used to select the candidates for multivariate analysis if the "p" value is < 0.05. The strength of association between dependent and independent variables (covariates) was expressed in odds ratio (OR). Finally, multivariate analysis using backward stepwise multiple logistic regression technique had been done to evaluate the independent effect of each variable on skilled attendance service utilization by controlling the effect of other confounders.

Operational Definitions

Bad obstetrics outcome: Those outcomes related with abortion, still birth, obstructed and prolonged labour.

Births in the last five years: All births within five years irrespective of outcome of delivery (live birth, abortion, stillbirth and death after live birth).

Know danger signs of labor: At least three of the accepted danger signs mentioned.

Know danger signs of pregnancy: At least three of the accepted danger signs mentioned.

Maternal Mortality: Is 'the death of a woman while pregnant or within 42 days of termination of pregnancy, irrespective of the duration and site of the pregnancy, from any cause related to or aggravated by the pregnancy or its management but not from accidental or incidental causes' [29].

Safe delivery: Normal labor and delivery attended in the health facilities by health care providers including health extension workers.

Skilled birth attendant: A medically qualified provider with midwifery skills (midwife, nurse or doctor) who has been trained to proficiency in the skills necessary to manage normal deliveries and diagnose, manage, or refer obstetric complications [28].

Traditional birth attendant (TBA): A community-based provider of care during pregnancy and childbirth. TBAs are not trained to proficiency in the skills necessary to manage or refer obstetric complications [28].

Results

Socio-demographic characteristics of the respondents

There are 500 women who had children less than 5 years old were participated in this study. Majority of participants 356 (71.2%) were in the age group of 26 - 30 years with Mean age of $27.44 \pm (3.076)$ years ranging from 20 - 37 years. Most of participants 454 (90.8%), 464 (92.8%) were Gamo by ethnicity and married respectively. Regarding religion more than one half 262 (52.4%) was protestant followed by orthodox 190 (38.0%). Concerning educational status one half 252 (50.4%) of mothers were illiterate and less than one fourth 116 (23.2%) of fathers were illiterate. With regard to Majority 336 (67.2%) of respondents have family size of 1 - 5 and Mean family size $4.87 \pm (1.315)$ Years, ranging from 3 - 9. More than one quarter 142 (28.4%) of respondents earn monthly income of less than 2000 ETB with Mean monthly income of $3121.04 \pm (SD = 1697.194)$ ETB, ranging from 1000 - 8600 (Table 1).

Socio-demographic characteristics		Frequency	Percent
Age category	15 - 20 years	7	1.4
	21 - 25 years	69	13.8
	26 - 30 years	356	71.2
	Above 30 years	68	13.6
Religion	Orthodox	190	38.0
	Protestant	262	52.4
	Muslim	22	4.4
	Catholic	26	5.2
Ethnicity	Gamo	454	90.8
	Gofa	15	3.0
	Amhara	17	3.4
	Wolayita	8	1.6
	Others	6	1.2
Marital status	Married	464	92.8
	Separated	29	5.8
	Divorced	7	1.4
Educational status of mother	Illiterate	252	50.4
	Completed primary school	117	23.4
	Completed secondary school	78	15.6
	College and University	53	10.6
Educational status of the father	Illiterate	116	23.2
	Completed primary school	134	26.8
	Completed secondary school	124	24.8
	College and University	126	25.2
Occupational status of Mother	Housewife	160	32.0
	Farmer	122	24.4
	Merchant	112	22.4
	Gov't employee	61	12.2
	Day laborer	18	3.6
	Private employee	24	4.8
	Others	3	.6
Occupational status of the husband	Farmer	196	39.2
	Merchant	151	30.2
	Government employee	74	14.8
	Private employee	48	9.6
	Daily Laborer	22	4.4
	Retired	9	1.8
Family size	1-5	336	67.2
	Above 5	164	32.8
Monthly in come in ETB	Less than 2000	142	28.4
	2001 - 3000	128	25.6
	3001 - 4000	95	19.0
	4001 - 5000	67	13.4
	Above 5000	68	13.6

Table 1: Frequency distribution of Socio-demographic characteristics of Mothers having children less than 5 years in Chencha District, Southern Ethiopia, December 2016 (n = 500).

Obstetric characteristics of the respondents

Majority 376 (75.2%) of women had married at age of 20 - 24 years. mean age of first marriage was $21.3020 \pm (2.18457)$ years, ranging from 17 - 32 years. Regarding Age at first pregnancy majority 390 (78.0%) and mean age at pregnancy was 21.6 ± 2.13192 Years, ranging from 17 - 32. With regard to outcomes of pregnancy most 469 (93.8%) had live birth followed by 14 (2.8%) abortion. Among mothers who were pregnant during study period majority 43 (79.6%) started ANC follow-up. Most of mothers 470 (94.0%) reported that their last pregnancy was planned and most of them reported that they had received ANC service for this pregnancy, however only about one third 147 (33.7%) had received ANC service for times. Majority 348 (69.6%) had delivered their last baby in home with most common mode of delivery being spontaneous vaginal delivery 457 (91.4%). Regarding knowledge and attitude of the mother 291 (58.2%) and 379 (75.8%) had adequate knowledge about institutional delivery and positive attitude about it respectively (Table 2).

Obstetric characteristics of the respondents	Frequency	Percent	
Age at marriage	14 - 19 years	101	20.2
	20 - 24 years	376	75.2
	25 - 30 years	23	4.6
Age at first pregnancy	14 - 19 years	81	16.2
	20 - 24 years	390	78.0
	25 - 34 years	29	5.8
Outcomes of pregnancy	Live birth	469	93.8
	Abortion	14	2.8
	Still birth	8	1.6
	Died within seven days	6	1.2
	Died b/n 7 days and birth	3	.6
Are you pregnant now	Yes	54	10.8
	No	446	89.2
How old is your pregnancy in months now (n = 54)	1 - 3 months	16	29.6
	4 - 6 months	32	59.3
	Above 6 months	6	11.1
Have you started ANC follow - up (n = 54)	Yes	43	79.6
	No	11	20.4
When was the last pregnancy in year	Less than one year	54	10.8
	1 year	338	67.6
	2 years	82	16.4
	3 years	20	4.0
	4 years	6	1.2
Was last pregnancy planned	Yes	470	94.0
	No	30	6.0
Did you receive ANC care	Yes	442	88.4
	No	58	11.6
Frequency of ANC visit (n = 442)	Once	43	9.7
	Twice	56	12.7
	Three times	194	43.9
	Four times	149	33.7
Place of delivery of last baby	Home	348	69.6
	Health Facility	152	30.4
Mode of last delivery	Spontaneous vaginal delivery	457	91.4
	Instrumental delivery	8	1.6
	Caesarean section	35	7.0
Who assisted you in last delivery	Physician	26	5.2
	Midwifery	106	21.2
	Nurse	20	4
	TBA	348	69.6
Condition of last baby	Live birth	468	93.6
	Live birth but died soon after birth	23	4.6
	Still birth	9	1.8
Did you faced Health problem faced During Labor and delivery	Yes	26	5.2
	No	474	94.8

Table 2: Frequency distribution of Obstetric characteristics of the respondents of Mothers having children less than 5 years in Chencha District, Southern Ethiopia, December 2016.

Regarding reported reasons of why mothers prefer home delivery majority 138 (39.7%) reported previous home delivery was normal, 121 (34.8%) reported labor was small and short, 36 (10.3%) they were told that there pregnancy was normal and 21 (6.0%) lack of transport and presence of traditional birth attendants (Figure 1-3).




Figure 1: Distribution of client related factors why mothers prefer home delivery in Chencha District, Southern Ethiopia, December 2016 (n = 348).




Figure 2: Distribution of health facility related factors why mothers prefer to give birth at home in Chencha District, Southern Ethiopia, December 2016 (n = 348).




Figure 3: Distribution of Reasons why mothers prefer give birth at health facility in Chencha District, Southern Ethiopia, December 2016 (n = 152).

Regarding health facility related factors why mothers prefer to give birth at home were 154 (44.3%) health facility is too far, 146 (42.0%) poor quality service, 32 (9.2%) no female provider at health facility and 16 (4.6%) unwelcoming approach of health care providers (Figure 4).




Figure 4: Health problem faced During Labor and immediately after delivery in Chencha District, Southern Ethiopia, December 2016 (n = 26).

Concerning reasons why mothers give birth at health facility more than one third 37.0% reported need better service, 28% reported previous better outcome in health facility delivery, 14% difficult labor and 7.0% bad outcome with previous home delivery (Figure 5).




Figure 5: Birth order of the last baby in Chencha District, Southern Ethiopia, December 2016 (n = 500).

Concerning Health problem faced During Labor and immediately after delivery the most common problem were 8 (30.8%), 6 (23.1%), 3 (11.5%) and 3 (11.5%) retained placenta, prolonged labor, mal-presentation and excessive vaginal bleeding respectively (Figure 6).




Figure 6: Conceptual frame work showing relationship between SBA utilization and associated factors reviewed from different literatures.

Regarding the birth order the last pregnancy of 193 (39.0%), 170 (34.0%), 118 (23.0%) and 19 (4.0%) of mothers was third, fourth and above, second and first respectively (Figure 7).




Figure 7: Schematic presentation of sampling.

Knowledge about pregnancy, labor and delivery service

Mothers were asked a series of questions regarding their knowledge about pregnancy, labor and delivery service and most of them 446 (89.2%) reported that they know health risks associated with pregnancy, more than one half 291 (58.2%) know danger signs of pregnancy, 295 (59.0%) know danger signs of labor, 387 (77.4%) reported having birth at home has associated risks and they also reported that having birth at health facility has associated benefit. Regarding risks associated with risks during pregnancy about one half 214 (48.0%) reported pregnancy related disease, 157 (35.2%) reported maternal death and 75 (16.8%) fetal death. Concerning reported danger signs leakage of amniotic fluid without labor 57 (19.6%), swelling of leg/face 46 (15.8%) followed by increased BP 37 (12.7%) and 34 (11.7%) severe headache and excessive weight gain. With regard to reported danger signs of labor, prolonged labor greater than 12 hrs 70 (23.7%), Increased BP 65 (22.0%) followed by convulsion 35 (11.9%) and Mal-presentation 35 (11.9%). Among reported benefits of giving birth at Health facility early detection of problems 89 (23.0%), low maternal mortality 69 (17.8%) followed by low maternal exhaustion 64 (16.5%). Overall knowledge of mother about pregnancy, labor and delivery service was adequate 291 (58.2%) and poor for 209 (41.8%) (Table 3).

Knowledge about pregnancy, labor and delivery service		Frequency	Percent
Awareness of health risk during pregnancy	Yes	446	89.2
	No	54	10.8
If yes what are risks (n = 446)	Pregnancy related disease	214	48.0
	Maternal Death	157	35.2
	Fetal death	75	16.8
Knowledge on danger signs of pregnancy	Yes	291	58.2
	No	209	41.8
If yes what are danger signs (n = 291)	Swelling of leg/face	46	15.8
	Vaginal bleeding	26	8.9
	Reduced fetal movement	28	9.6
	Severe headache	34	11.7
	Severe abdominal cramp	29	10.0
	Excessive weight gain	34	11.7
	Increased BP	37	12.7
	Leakage of amniotic fluid without labor	57	19.6
Do you know danger signs of labor	Yes	295	59.0
	No	205	41.0
If yes what are danger signs of labor (n = 295)	Prolonged labor greater than 12hrs	70	23.7
	Early rupture of membrane	25	8.5
	Vaginal bleeding	20	6.8
	Placental retention greater than 1hr	10	3.4
	Mal-presentation	35	11.9
	Increased BP	65	22.0
	Convulsion	35	11.9
	Abdominal pain	35	11.9
Giving birth at home has risks	Yes	387	77.4
	No	113	22.6
If yes what do you know (n = 387)	Maternal exhaustion	64	16.5
	Fetal distress	113	29.2
	Maternal death	89	23.0
	Fetal/neonatal death	58	15.0
	Postpartum morbidity	41	10.6
	Disease transmission from attendants	22	5.7
Is there any benefit in giving birth at Health facility	Yes	387	77.4
	No	113	22.6
If yes what are they (n = 387)	Early detection of problems	89	23.0
	Early treatment of problems	47	12.1
	Lower maternal exhaustion	64	16.5
	Better newborn care	43	11.1
	Lower maternal mortality	69	17.8
	Low postpartum morbidity	75	19.4
Overall Knowledge on pregnancy, labor and delivery service	Adequate Knowledge	291	58.2
	Poor Knowledge	209	41.8

Table 3: Frequency distribution of Mothers Knowledge about pregnancy, labor and delivery service in Chencha District, Southern Ethiopia, December 2016 (n = 500).

Knowledge: Labelled to have adequate Knowledge if they correctly responded all questions designed to access knowledge about pregnancy, labor and delivery.

Attitude about pregnancy, labor and delivery service

Mothers were asked series of questions concerning their perception about pregnancy, labor and delivery complications and majority reported that complication of labor were preventable and treatable 380 (76.0%) and 439 (87.9%) respectively. Most of respondents 400 (80.0%) reported that every woman was at risk of developing pregnancy complication. Overall attitude of mothers about pregnancy, labor and delivery was positive for 379 (75.8%) and negative for 121 (24.2%) (Table 4).

Attitude about pregnancy, labor and delivery service	Attitude score	
	Positive (1)	Negative (0)
Do you think most complications of labor were preventable	380 (76.0%)	120 (24.0%)
Do you think most complications of labor were treatable	439 (87.8%)	61 (12.2%)
Any pregnant women are at risk for developing delivery complications	400 (80.0%)	100 (20.0%)
Delivery complications are hazardous to maternal health	458 (91.6%)	42 (8.4%)
Delivery complications are risk for newborn	412 (82.4%)	88 (17.6%)
Skilled birth attendance is important for my wellbeing	434 (86.8%)	66 (13.2%)
Skilled birth attendance is important for new born baby	382 (76.4%)	118 (23.6%)

Table 4: Frequency distribution of Mothers Attitude about pregnancy, labor and delivery service in Chencha District, Southern Ethiopia, December 2016 (n = 500).

Note: Very agree and Agree were labeled as positive and assigned value "1" and Disagree and very disagree were labeled as negative attitude and assigned value "0". The Overall Attitude was considered positive if all questions were assigned value "1" and respondents were labeled to negative attitude if any of questions is assigned "0".

Social and Health system factors and women decision-making

Most of mothers reported that they decide on utilization of health service and place of delivery 349 (69.8%) and 312 (62.4%) respectively. Concerning health system related factors majority of respondents 363 (72.6%), 337 (67.4%), 320 (64.0%) and 180 (36.0%) reported that Health professionals are skilled enough, The facility is well equipped to provide service, distance from health facility was greater than 5km and nearby health facility was health center respectively (Table 5).

Social factors		Frequency	Percent
Who decides your health service utilization	Self	349	69.8
	Husband	137	27.4
	Relatives	14	2.8
Who decides your place of childbirth	Self	312	62.4
	Husband	132	26.4
	Relatives	43	8.6
	Religious leader	13	2.6
Health system factors			
Health professionals are skilled enough	Yes	363	72.6
	No	137	27.4
The facility is well equipped to provide service	Yes	337	67.4
	No	163	32.6
Distance of nearby Health facility in Km	Less than 5 km	180	36.0
	Greater than 5 km	320	64.0
Types of nearby health facility	Health post	160	32.0
	Hospital	160	32.0
	Health center	180	36.0

Table 5: Frequency distribution of social and system factors associated with pregnancy, labor and delivery service in Chencha District, Southern Ethiopia, December 2016 (n = 500).

Factors associated with skilled birth attendance

Binary logistic regression was done to identify association between skilled birth attendance utilization and independent variables and variables that have shown statistical significance at P-value less than 0.05 were entered into multinomial logistic regression to rule out confounders and identify independent predictors of skilled birth attendance utilization practice. Age of the mother 21 - 25 years were 1.5 times [AOR = 1.56 (1.204 - 3.516)] more likely to deliver their babies in health facility than those above 30 years of age, Mothers with government employed husband were 1.6 times [AOR = 1.669 (1.475 - 81.642)] more likely more likely to deliver their babies in health facility than farmers, mothers with monthly income above 5000 ETB were 2 times [AOR = 2.196 (1.051 - 4.590)] more likely to deliver their babies in health facility than those earning below 200 ETB per month, Mothers with no ANC follow-up were less [AOR = 0.915 (0.375 - 0.953)] likely to deliver in health facility than those who have ANC follow-up, Mothers with four times ANC visit were 5 times more [AOR = 5.615 (1.851 - 17.040)] likely to deliver in health facility than those with one time follow-up, Mothers who reported health professionals

are not well skilled were less [AOR = 0.227 (0.085 - 0.607)] likely to deliver in health facility than those who reported professionals are well skilled, Mothers with nearest health facility in less than 5 km distance were 3 times more [AOR = 3.116 (1.916 - 5.068)] likely to deliver in health facility than those with more than 5km distance (Table 6).

Modern contraceptive use ^a Home (n = 348) Frequency Percent		Place of delivery of last baby in 5yrs				COR UB	95% CI for AOR		AOR UB	95% for AOR		P value ^b	
		Health Facility (n = 152)		LB			LB	LB		LB	LB		
		Frequency	Percent										
Home	Age of the respondent	15-20 years	5	1.4	2	1.3	1.30	.230	7.354	.600	.083	4.322	.767
		21-25 years	39	11.2	30	19.7	2.50	1.199	5.215	1.56	1.204	3.516	.015*
		26-30 years	252	72.4	104	68.4	1.341	.732	2.456	.772	.348	1.712	.342
		Above 30 years	52	14.9	16	10.5	.947	.229	3.911				
	Occupation of the husband	Farmer	133	38.2	63	41.4	1	-	-	1	-	-	-
		Merchant	101	29.0	50	32.9	1.364	.316	5.881	.719	.070	7.361	.989
		Government employee	44	12.6	30	19.7	.133	.022	.817	1.669	1.475	81.642	.019*
		Private employee	45	12.9	3	2.0	.316	.050	1.998	6.226	.064	7.013	.677
		Daily Laborer	19	5.5	3	2.0	.947	.229	3.911	1.760	.133	23.339	.221
		Retired	6	1.7	3	2.0	.990	.238	4.124	.534	.053	5.378	.940
	Monthly income	Less 2000 ETB	103	29.6	39	25.7	1	-	-	1	-	-	-
		1000-3000 birr	86	24.7	42	27.6	.657	.358	1.204	1.448	.689	3.042	.174
		3001-4000 birr	74	21.3	21	13.8	.382	.193	.755	1.846	.833	4.088	.006
		4001-5000 birr	46	13.2	21	13.8	.614	.303	1.243	1.196	.515	2.779	.175
		Above 5000 birr	39	11.2	29	19.1	.509	.278	.933	2.196	1.051	4.590	.029*
	Did you receive ANC care	Yes	316	90.8	126	82.9	1	-	-	1	-	-	-
		No	32	9.2	26	17.1	.491	.281	.857	.915	.375	.953	.012
	Frequency of ANC visit	Once	29	8.3	14	9.2	1	-	-	1	-	-	-
		Twice	51	14.7	5	3.3	.901	.438	1.853	.915	.375	2.233	.845
		Three times	139	39.9	55	36.2	.738	.466	1.168	1.340	.762	2.356	.309
Four times		97	27.9	52	34.2	.183	.069	.486	5.615	1.851	17.040	.002*	
Health professionals are skilled enough	Yes	238	68.4	125	82.2	1	-	-	1	-	-	-	
	No	110	31.6	27	17.8	.160	.039	.651	.227	.085	.607	.003*	
Distance from nearest health facility	Less than 5 km	145	41.7	35	23.0	.419	.271	.646	3.116	1.916	5.068	0.000*	
	Greater than 5 km	203	58.3	117	77.0	1	-	-	1	-	-	-	

Table 6: Multivariable logistic regression of factors predicting the likelihood of modern family planning method utilization among reproductive age women at Arba Minch town, Southern Ethiopia, 2017, (n = 500).

^a95% Confidence Interval for place delivery

^bP-value between groups significant at the 0.05 level

Discussion

This study revealed the status and independent predictors of skilled birth attendance by rural mothers in Chencha district, Gamo Gofa zone southern Ethiopia. The point prevalence of institutional delivery was 152 (30.4%) and Majority 348 (69.6%) had delivered their last baby in home with most common mode of delivery being spontaneous vaginal delivery 457 (91.4%). This is higher than study conducted in southern Ethiopia showed that the prevalence of skilled birth attendance was 148 (26.8%) [22] and findings from other rural communities Ethiopia For example 21% of births to rural mothers were assisted by a skilled provider and 20% were delivered in a health facility [6]. However this is lower than study conducted in North West Ethiopia showed that 38.2% of women gave birth in health facilities for their most recent birth while the most 223 (61.8%) delivered at home [14]. The variation could be explained by difference in socio-demographic variables among study participants and difference in tools used by researchers.

This study showed that mothers knowledge about pregnancy, labour and delivery was adequate for the 291(58.2%) of them. However knowledge of mothers was not associated with skilled birth attendance. This could be due others factors deterring the use skilled birth attendance like maternal age, monthly income, occupation, skill of professionals and distance from health facility.

This study revealed that the major reasons why mothers give birth at health facility include; 37.0% need better service, 28% previous better outcome in health facility delivery, 14% difficult labor and 7.0% bad outcome with previous home delivery. This is similar to findings from north west Ethiopia showed that major reasons of institutional delivery were fear of complications (77.5%), to get better service (56.5%) safe and clean delivery (46.4%) and they were communicated to deliver at health facilities during ANC attendance, 47 (35.5%) [14].

Age of the mother 21 - 25 years were 1.5 times [AOR = 1.56 (1.204 - 3.516) more likely to deliver their babies in health facility than those above 30 years of age. This in line with findings from different studies conducted in north west Ethiopia indicated that mothers with lower age at first pregnancy, having more than one previous pregnancies, with in more than one birth order, received ANC service, found to be significantly associated with safe delivery service utilization [18]. Study conducted in southern Ethiopia showed that Mothers with the age range of 15-19 yrs were about five times more likely to give birth at health facility [22]. This could be due to improved access to reproductive health information for young generation through Medias like, TV, radio or internet than those old aged mothers.

Mothers with government employed husband were 1.6 times [AOR = 1.669 (1.475 - 81.642) more likely more likely to deliver their babies in health facility than farmers. This is in line with studies conducted in different regions of Ethiopia indicated that maternal education family size, husband occupation and education were all significantly associated with health facility delivery [16].

Mothers with monthly income above 5000 ETB were 2 times [AOR = 2.196 (1.051 - 4.590) more likely to deliver their babies in health facility than those earning below 200 ETB per month. This is in line with EDHS findings household wealth status is highly correlated with whether their deliveries are assisted by a skilled provider and whether they are delivered in a health facility [6]. This could be due to increased wealth of family could help other factors hindering institutional delivery like distance from health facility by that they can pay transportation cost to get ANC service or attend delivery in health facility.

Mothers with no ANC follow-up were less [AOR = 0.915 (0.375 - 0.953) likely to deliver in health facility than those who have ANC follow-up, Mothers with four times ANC visit were 5 times more [AOR = 5.615 (1.851 - 17.040) likely to deliver in health facility than those with one time follow-up. This similar with findings from Study conducted in North West Ethiopia showed that mothers who received ANC service, birth complications in previous and immediate pregnancies and had every used professionally assisted delivery in preceding delivery were commonly found to be significantly associated with safe delivery service utilization [18]. Antenatal care service utilization is independent predictor of utilization of institutional delivery. Mothers making one or more ANC visits were more likely to use delivery care service. This might be due to advice from healthcare workers during antenatal care increases a woman's use of institutional delivery. The

healthcare workers could provide good information regarding safe health care delivery and encourage women to deliver at health facility. A previous study in Ethiopia found that proper counseling and advice to deliver at health care facility increased institutional delivery [19].

Mothers who reported health professionals are not well skilled were less [AOR = 0.227 (0.085 - 0.607)] likely to deliver in health facility than those who reported professionals are well skilled. This is similar with study in Cambodia indicated that delivery with skilled attendant at the preceding delivery was a significant determinant for subsequent use of skilled attendant. Once a woman has delivered with the aid of an unskilled attendant, she is five to seven times less likely to seek skilled help than a primipara [20].

Mothers with nearest health facility in less than 5 km distance were 3 times more [AOR = 3.116 (1.916-5.068)] likely to deliver in health facility than those with more than 5km distance. This similar with findings from Bahir Dar, showed that Mothers who gave birth at home have different reasons 46% of them think giving birth at health facility has no necessity, 33% of mothers said the service at health facility is not customary, 21% of the mother said that the health facility was either too far or that they did not have transportation. Rural women were more likely to report that facility deliveries are not customary 33%, health facilities were too far or they had no transportation 22% [23].

Conclusion

In conclusion this study revealed that prevalence of skilled birth attendance below one third. ANC visit for four and more times low, Mothers knowledge about pregnancy, labour and delivery was low. Age of the mother 21 - 25 years, Mothers with government employed husband, mothers with monthly income above 5000 ETB, Mothers with ANC follow-up, Mothers with four times ANC visit, Mothers who reported health professionals are not well skilled and Mothers with nearest health facility in less than 5 km distance were predictors of skilled birth attendance utilization.

Strengths and Limitations of the Study

Strengths

Random sampling technique was employed. i.e., random selection of the Villages, Data collectors speaks local languages and was the same sex with study subjects to minimize information biases and Standardized questionnaire based on similar studies.

Limitations

The study design was cross-sectional and determinants of skilled birth attendance cannot assessed, there could be recall bias, since the women were asked for events within the last one year prior to the survey.

Recommendations

Based on the findings of this study, the following specific recommendations were made to respective bodies.

To health care providers

- The knowledge of mothers regarding pregnancy, labour and delivery was low; this recalls Health care providers should provide information on risks of pregnancy, benefits of giving birth at health facilities, risk of giving birth at home to mothers, family members and the community.
- Mothers who had ANC follow-up four and more times were still low professionals should promote it to four and more and linked with delivery care.
- Providing standardized care for clients is the basic concern for every professional and it is important to update skill and knowledge to provide care.

To woreda health sector officials

- Improving access and quality of health service were independent predictors of skilled birth attendance. Therefore availing basic supplies and staffing with trained health professionals)
- More work is expected to achieve the national need i.e. creating home delivery free village and preventing all preventable maternal deaths in 2030.

To community

1. Community leaders should encourage health facility visit during pregnancy, labour and delivery and discourage involvement of traditional birth attendants on delivery assistances.

Contribution of Researcher

Mende Mensa is senior researchers who conceived the study and prepared the proposal; analyzed the data and presented the work for responsible bodies analyzed and interpreted the findings of this study and he also prepared this document for publication. Melkenesh Nigatu has participated in data entry and analysis.

Conflicts of Interest

We have no conflict of interest during conducting this study or developing the manuscript. All expenses regarding this paper were from out of pocket of the researchers.

Acknowledgements

Above all our thanks go to Almighty God, Who provided us time, health and endurance to conduct this study. We would like to express our deepest gratitude to our friends Mr. Habitamu Samuel (Health officer, Coordinator of research and publication core process owner), for his unreserved help during analysis, interpretation of statistical data and the writing of final paper. Our appreciation also goes to all staffs of Arba Minch college health Sciences for their unreserved support throughout the course and final thesis writing. Our heartfelt gratitude and thanks to staffs of Gamo Gofa Zone health department, Chench district council and health office and mothers participated in this study for offering us all necessary materials and vital information during data collection and writing final thesis . Really, we are very grateful for your whole rounded support. Thank you very much! We are heartily thankful to our children, Kalkidan and Mikiyas and all other family members and friends for their continuous encouragement, moral and material support throughout this research work. Your closeness was very vital for our success. Really, I am proud of you!

Ethical Consideration

Ethical clearance was obtained from institutional Review Board of Arba Minch College of Health Sciences. Permission letters to conduct the study was obtained from south nation nationalities and people region health bureau, Gamo Gofa zone health department, Chench woreda health office and respective kebele administrations. Aim of the study was explained, informed written consent had been obtained from each participant. Confidentiality was maintained by using codes that is only known by the respondent. Participants who were unwilling to participate in the study and those who wanted to abstain from their participation at any stage were informed to do so without any restriction.

Bibliography

1. World Bank. Population and reproductive Health, World Bank (2002).
2. Ethiopian Society of Population Studies. "Maternal Healthcare Seeking Behavior in Ethiopia: Findings from EDHS 2005" (2008).

3. Economic Commission For Africa African Union Bank Africa Development Programme, and United Nation Development Programme. Millennium development goal report 2012: Assessing progress in Africa toward the millennium development goals, Emerging perspectives from Africa on the post-2015 development agenda.
4. WHO. "Making pregnancy safer: the critical role of the skilled attendant". Department of Reproductive Health and Research, Geneva (2004).
5. WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division. Trends in maternal mortality: 1990 to 2015 (2015).
6. Central statistics Agency. Ethiopia demographic and health survey 2016, Addis Ababa, Ethiopia (2016).
7. Making pregnancy safer: the critical role of the skilled attendant: a joint statement by WHO, ICM and FIGO. Geneva: World Health Organization (2014).
8. Lawn JE., et al. "Stillbirths: Where? When? Why? How to make the data count?" *Lancet* 377.9775 (2011): 1448-1463.
9. Montagu D., et al. "Where do poor women in developing countries give birth? A multi-country analysis of demographic and health survey data". *PLoS One* 6 (2011): e17155.
10. UN. End Poverty 2015 makes it happen. Millennium Development Goals Fact Sheet. United Nation Headquarter, New York (2008).
11. Graham W., et al. "Can skilled attendances at delivery reduce maternal mortality in developing countries?" *Studies in Health Services Organisation and Policy* 17 (2001): 97-129.
12. WHO (1990) UNFPA, World Bank Group, United Nations Population Division. Trends in maternal mortality, 2015.
13. Oestergaard MZ., et al. "Neonatal mortality levels for 193 countries in 2009 with trends since 1990: a systematic analysis of progress, projections, and priorities". *PLoS Medicine* 8.8 (2011): e1001080.
14. Belay and Sendo. "Factors determining choice of delivery place among women of child bearing age in Dega Damot District, North West of Ethiopia: a community based cross-sectional study". *BMC Pregnancy and Childbirth* 16 (2016): 229.
15. Thaddeus S and Maine D. "Too far to walk: Maternal mortality in context: Findings from a multidisciplinary literature review. Columbia University School of Public Health: Prevention of Maternal Mortality Program, Center for Population and Family Health (1990).
16. Abyot Asres. "Assessment of factors associated with safe delivery service utilization among women of child bearing age in Sheka zone, snnpr, south west Ethiopia". Addis Ababa University Faculty of Medicine School of Public health (2008).
17. FL Fredrik and G Langi GL. "Barriers to delivery care by skilled attendants in Sulawesi Utara". *Journal of Public Health and Development* 7.3 (2009): 1-13.
18. Pfeiffer C and Mwaipopo R. "Delivering at home or in a health facility? Health seeking behavior of women and the role of traditional birth attendants in Tanzania". *BMC Pregnancy Childbirth* 13.1 (2013): 55.
19. Mesfin Nigussie., et al. "Assessment of safe delivery service utilization among women of childbearing age in north Gondar Zone, north west Ethiopia". *Ethiopian Journal of Health Development* 18.3 (2004): 145-152.
20. Yanagisawa S., et al. "Determinants of skilled birth attendance in rural Cambodia". *Tropical Medicine and International Health* 11.2 (2006): 238-251.
21. Yifru B. "A metaanalysis of sociodemographic factors predicting birth in health facility". *Ethiopian Journal of Health Sciences* 24 (2014): 81-92.

22. Kaleb Mayisso Rodamo., *et al.* "Magnitude and Determinants of Utilization of Skilled Birth Attendance among Women of Child Bearing Age in Sidama Zone, Southeast Ethiopia". *Journal of Gynecology and Obstetrics* 3.4 (2015): 69-76.
23. Abeje G., *et al.* "Factors associated with Institutional delivery service utilization among mothers in Bahir Dar City administration, Amhara region: A community based cross sectional study". *Reproductive Health* 11.1 (2014): 22.
24. Hailu M., *et al.* "Birth preparedness and complication readiness among pregnant women in Southern Ethiopia". *PLoS One* 6.6 (2011): e21432.
25. CSA [Ethiopia]. Ethiopia Mini Demographic and Health Survey 2014. Addis Ababa, Ethiopia (2014).
26. World Health Organization. WHO 2012 Maternal and Child Health Fact Sheet (2012).
27. Zelalem T., *et al.* "Magnitude and Factors Associated with Institutional Delivery Service among Women who have Antenatal Care Follow-Up at Hawassa University Referral Hospital". *Diversity and Equality in Health and Care* 14.1 (2017): 16-22.
28. WHO Maternal and Newborn survival and health guidelines (2006), Integrating management of pregnancy and childbirth (WHO 2007) and UNFPA Safe Motherhood program (2007).
29. WHO. 'Maternal Mortality Ratio'. Health statistics and health information systems (2011).

Volume 5 Issue 6 September 2017

© All rights reserved by Mende Mensa and Melkenesh Nigatu Belijo.