

## **Golden Age of Green Chemistry**

## Ana Carolina Kogawa and Hérida Regina Nunes Salgado\*

São Paulo State University (UNESP), School of Pharmaceutical Sciences, Araraquara, SP, Brazil

\*Corresponding Author: Hérida Regina Nunes Salgado, Professor, São Paulo State University (UNESP), School of Pharmaceutical Sciences, Araraquara, SP, Brazil.

Received: September 21, 2017; Published: September 22, 2017

This is the golden age of Green Chemistry. Many analytical optimizations have made relevant differences to the community, as well as the chemical industries and the environment. The presence of pharmaceutical residues or chemical solvents in the environment has become a recent research issue. In fact, tons of them are annually produced around both the developing and developed countries. In this scenario we have to think about it. Could we be done research and the improvements better? In face of pressures to reduce the amount of residues produced, the researchers carry out green techniques.

The human and environmental contamination of chemical substances has been noticed in academic and social websites and consequently compelling these scientific committees as a potential topic meriting concern.

Pharmaceutical and chemical industries obviously differ concerning the residues produced or the amount of residues. Apart from the potential for soil contamination and consequently groundwater, the waste reaches first the man or the analyst who has direct contact every day with solvents and reagents. This affects the health and consequently quality of life of them. Residues contaminate plantations, fish and animals and are further disseminated through the food chain. Green chemistry comes to avoid neglect with the operator, animals, vegetation and everything that lives on our planet. It comes to bring quality!

Analytical methods aims to propose a rapid, sensitive, easy to perform and reproducible procedure able to quantify the active pharmaceutical ingredients (API) in both raw material and pharmaceutical products [1-3]. An example of a green method is the infrared (IR) spectrophotometry.

IR is a relatively new and ecological quantitative technique that makes the resolution possible in significantly less residue production because of its very small quantity of KBr spend (approx. mg) with significantly lower consumption of material. Usually infrared spectrophotometric analysis are conducted to qualitative characterization of drugs, such can be accessed in many pharmacopeias or other official compendia. However, the infrared spectrophotometry could be studied in detail, in order to exactly quantify drugs and pharmaceuticals, such can be found in the literature [4-18]. These methods aimed the quantification of pharmaceuticals quickly, cheaply and without toxic solvents. This contemplates the thought of ecologically correct method.

In the near future, the validation of green methods should be set out as a priority for the academy as well as the general society. Additionally, it is very important to validate new analytical methods and divulgate these protocols around the world in order to protect the future generations, animals and our planet. This is the future for chemical-pharmaceutical analyzes. The one-dimensional thinking of just validating methods to obtain an effective result has become outdated. Currently the thought must be multi-dimensional; methods must be validated to get an effective, cheap, fast and green result [19].

## **Bibliography**

1. Bonfilio R., *et al.* "Recent applications of analytical techniques for quantitative pharmaceutical analysis: a review". *WSEAS Transac*tions on Biology and Biomedicine 7.4 (2010): 316-338.

Citation: Ana Carolina Kogawa and Hérida Regina Nunes Salgado. "Golden Age of Green Chemistry". EC Microbiology 12.2 (2017): 52-54.

- 2. Bonfilio R., *et al.* "Analytical validation of quantitative high-performance liquid chromatographic methods in pharmaceutical analysis: A practical approach". *Critical Reviews in Analytical Chemistry* 42.1 (2012): 87-100.
- 3. Kogawa AC and Salgado HRN. "Analytical methods need optimization to get innovative and continuous processes for future pharmaceuticals". Scholars Academic Journal of Pharmacy 5.6 (2016): 240-244.
- 4. Matkovic SR., *et al.* "Desarrollo y validación del análisis cuantitativo de ibuprofenoen comprimidos por espectroscopia infrarroja". *Acta Farmacéutica Bonaerense* 23.4 (2004): 527-532.
- Farouk F., et al. "Fourier transform infrared spectroscopy for in-process inspection, counterfeit detection and quality control of antidiabetic drugs". Spectroscopy 26.4-5 (2011): 297-309.
- 6. Sherazi STH., *et al.* "Application of Fourier-transform infrared (FT-IR) transmission spectroscopy for the estimation of roxithromycin in pharmaceutical formulations". *Vibrational Spectroscopy* 55.1 (2011): 115-118.
- 7. Moreno AH and Salgado HRN. "Development and validation of quantitative analysis of ceftazidime in powder for injection by infrared spectroscopy". *Physical Chemistry* 2.1 (2012): 6-11.
- 8. Tótoli EG and Salgado HRN. "Development and validation of the quantitative analysis of ampicillin sodium in powder for injection by Fourier-transform infrared spectroscopy (FT-IR)". *Physical Chemistry* 2.6 (2012): 103-108.
- 9. Vieira DCM., *et al.* "Development and validation of the quantitative analysis of cefuroxime sodium in powder for injection by infrared spectroscopy". *Advances in Analytical Chemistry* 2.6 (2012): 80-87.
- 10. Tótoli EG and Salgado HRN. "Fourier-Transform Infrared (FT-IR) spectrophotometry: an eco-friendly method for the analysis of daptomycin injectable". *Journal of AOAC International* 100.5 (2017): 1569-1576.
- 11. Corrêa JCR and Salgado HRN. "A platform for designing quantitative infrared spectrophotometric method for drugs and pharmaceuticals analysis: a rediscover for an ecological and safer technique in the routine quality control laboratories". *World Journal of Pharmacy and Pharmaceutical Sciences* 3.6 (2014): 2056-2059.
- 12. Natori JSH., *et al.* "Development and validation of a green analytical method for determination of norfloxacin in raw material by Fourier-Transform Infrared Spectrophotometry (FT-IR)". *Journal of AOAC International* 99.6 (2016): 1533-1536.
- 13. Consortti LP and Salgado HRN. "Green method for quantification of sodium cefotaxime in lyophilized powder by infrared spectroscopy". *Journal of Pharmaceutical Sciences and Emerging Drugs* 5 (2017): 1-6.
- 14. Trindade MT and Salgado HRN. "Development and validation of a modern and stability-indicating method for the quantification of ceftriaxona sodium in powder for injection by infrared spectroscopy". *Physical Chemistry* 7.3 (2017): 55-62.
- 15. Figueiredo AL and Salgado HRN. "Validation of a green analytical method for the quantitative analysis of antimicrobial aztreonam in lyophilized powder for injection by Fourier-transform infrared spectroscopy (FT-IR)". *EC Microbiology* 8.5 (2017): 254-265.
- 16. Kogawa AC and Salgado HRN. "Spectrophotometry in infrared region: a new, low cost and green way to analyze tablets of rifaximin". *Current Pharmaceutical Analysis* (2017).
- 17. Kogawa AC and Salgado HRN. "Development and validation of infrared spectroscopy method for the determination of darunavir in tablets". *Physical Chemistry* 3.1 (2013): 1-6.

Citation: Ana Carolina Kogawa and Hérida Regina Nunes Salgado. "Golden Age of Green Chemistry". EC Microbiology 12.2 (2017): 52-54.

- 18. Kogawa AC., *et al.* "Quantification of doxycycline in raw material by an eco-friendly method of infrared spectroscopy". *Pharmaceutica Analytica Acta* 7 (2016): 463-466.
- 19. Kogawa AC and Salgado HRN. "Quality tools for successful strategic management". *International Journal of Business Process Integration and Management* 8.3 (2017): 153-159.

## Volume 12 Issue 2 September 2017 © All rights reserved by Ana Carolina Kogawa and Hérida Regina Nunes Salgado.