

## Distribution and Pathogenicity Profile of Dematiaceous Fungi Isolated at Uli Community, Anambra State

Okereke Ogonna Friday<sup>1</sup>, Osuji Malachy Ikeokwu<sup>1\*</sup>, Umedum Chinelo Ursula<sup>2</sup>, Unaegbu Valentine Nnachetam<sup>3</sup> and Nkechinyere Opara-Nadi<sup>1</sup>

<sup>1</sup>Department of Biological Sciences (Microbiology), Faculty of Natural and Applied Sciences, Spiritan University, Nneochi, Abia State, Nigeria

<sup>2</sup>Department of Medical Laboratory, Faculty of Health Sciences, Chukwumeka Odumegwu Ojukwu University, Igbariam Campus, Anambra State, Nigeria

<sup>3</sup>Department of Biological Sciences, University of Agriculture and Environmental Sciences, Umuagwo, Imo State, Nigeria

**\*Corresponding Author:** Osuji Malachy Ikeokwu, Department of Biological Sciences (Microbiology), Faculty of Natural and Applied Sciences, Spiritan University, Nneochi, Abia State, Nigeria.

**Received:** September 08, 2025; **Published:** January 29, 2026

### Abstract

Superficial mycosis has been described as the major source of debilitating skin infection, especially in rural areas. This study examined the distribution and pathogenicity profile of dematiaceous fungi isolated at Uli community, Anambra State. Two hundred and ten (210) soil samples were randomly collected from three different soil types (loamy, clay, and sandy soil) at Uli community using a soil auger. The soil samples were analyzed using standard microbiological technique. The fungal isolates were characterized based on their morphology, slide culture technique, and atlas of clinical mycology. The pathogenicity potential of the isolates was evaluated using Wistar rats. *Exophiala jeanselmei* was mostly isolated (40%) from Umuoma community, followed by *Cladophialophora carriponii* (30%) from the same community, while the least was *Scedosporium apiospermum* (10%) from Aluoha community. Loamy soil yielded the highest fungal isolates 13 (65%), followed by clay soil 4 (20%) while sandy soil yielded the least fungal isolates 3 (15%). Statistically, there was a significant difference ( $P < 0.05$ ) in the distribution of the fungal isolates in the different soil types. Umuoma yielded the highest number of fungal isolates 10 (50%), followed by Umuaku 6(30%) while the least was Aluoha 4 (20%). The Wistar rats infected using the isolates developed erythematous lesions, which were confirmed by culturing scrapings from the infected site. The study showed that dematiaceous fungi are mostly found in loamy soil, of which *Exophiala jeanselmei* had the most frequent occurrence. Also, dematiaceous fungal isolates were able to cause erythematous lesions on the laboratory animals, which confirmed their pathogenicity potentials.

**Keywords:** Dematiaceous Fungi; Erythematous Lesions; Infection; Pathogenicity

### Introduction

Dematiaceous fungi are group of organisms that are diverse in nature, having the ability to produce dark pigmentation known as melanin in their hyphae or spores [1]. Dematiaceous or dark fungi are highly saprophytic, and are capable of causing infections in humans, animals, and plants [2]. They can be grown on different microbiological media (Sabouraud dextrose agar, Potato dextrose agar), where they produce dark grey colonies, white, brown, and black on agar plate surface while reverse pigmentation is dark coloured. The melanin pigment has been described as a virulent factor [2]. Dark fungi are abundant in nature, some common genera include: *Cladosporium*, *Curvularia*, *Exophiala*, *Bipolaris* etc.

**Citation:** Osuji Malachy Ikeokwu., et al. "Distribution and Pathogenicity Profile of Dematiaceous Fungi Isolated at Uli Community, Anambra State". *EC Microbiology* 22.2 (2026): 01-05.

Research had shown that distribution of dark fungi is localized due to their saprophytic nature [2]. The organisms had been reported to inhabit environment that provides opportunity for degradation, due to their saprophytic nature [1]. Most of the fungi had been isolated in the soil, rotten wood, air, and water [1]. However, it is worthy to note that soil harbors most of the organisms, and their distribution in the soil is influenced by several factors such as soil structure and texture [3].

Dematiaceous fungi had been revealed to be responsible for subcutaneous infections in both humans and animals [1]. They are highly pathogenic to humans, both individuals that have vibrant immune defense mechanisms and immunodeficient are infected [2]. Individuals in rural areas that are engaged in farming activities, hunting, fetching of firewood are highly predisposed to the fungal agents, as they can enter the body through piercing or traumatic inoculation via broken wood or stick in the soil. The fungi usually attack the lower limbs with appearance of lesions [2]. These fungal species also disrupt the functioning of transplanted organ, resulting in morbidity as reported by Yew, *et al.* [2].

Several researchers had worked on the distribution of dematiaceous fungi in soil and their pathogenicity potential such as Yew, *et al.* [2], Revanker and Suton [1] and Chukwuma, *et al.* [3] but few research had been geared towards the distribution and pathogenicity profile of dematiaceous fungi isolated at Uli community. Hence, this research is aimed at evaluating the distribution and pathogenicity profile of dematiaceous fungi isolated at Uli community. The result obtained from this study would contribute immensely to checking the spread of fungal infection associated with dark fungi.

## Materials and Methods

### Collection of samples

Soil samples comprising of 70 loamy, 70 clay, and 70 sandy soil were collected from three communities (Aluoha, Umuaku, and Umuoma) at Uli, Ihiala L.G.A, Anambra State. The samples were collected using a sterile spoon at a depth of 10 cm. The samples were put in a sterile polythene bag and were conveyed to the Department of Microbiology Laboratory, COOU for analysis, which was carried out within 2h.

### Processing of samples

The soil samples were serially diluted using normal saline. Ten millimeter of normal saline was put into test tubes containing 1g of soil samples and tenfold serial was done to obtain the following dilutions:  $10^{-1}$ ,  $10^{-2}$ ,  $10^{-3}$ ,  $10^{-4}$ ,  $10^{-5}$ , and  $10^{-6}$ . Test tubes containing  $10^{-4}$  and  $10^{-5}$  dilutions were plated on Potato Dextrose Agar (PDA) using spread plate technique. All the inoculated plates in triplicates were incubated in inverted position at  $30 \pm 2^\circ\text{C}$  for 7 days. The isolated colonies were then purified using a freshly prepared PDA.

### Identification of the isolates

The fungal growths were thoroughly scrutinized morphologically and microscopically. In morphological characterization, the colors, texture, size, margin, and reverse pigmentation were examined. Further identification was done using slide culture technique which enabled the structure of the melanized fungi to be clearly visualized as elucidated by Umedium and Iheukwumere [4]. The shape, color, and size of the conidia and hyphae were examined with the aid of a digital microscope and the overall identification was made using color atlas of clinical mycology [4].

### Pathogenicity test

Two months Wistar rats numbering 18 with average weight of 150g were purchased, and acclimatized for 7 days. During this period, the rats were provided with water and growers' feed inside a metallic cage that had optimum ventilation. The inoculum of the isolates was prepared and standardized using 0.5 MacFarland solution. The Wistar rats were divided into four groups. Each of the isolates (0.1 mL) was inoculated into the rats subcutaneously using 1 mL syringe after disinfection and depilation using 70% ethanol and sterile blade, respectively. Dimethyl Sulfoxide (0.1 mL) was inoculated into the control rats. The infected rats were fed and observed for 21 days.


Erythematous lesions that developed on the infected site on the 14<sup>th</sup> day were scrapped and cultured on SDA and incubation followed at  $30 \pm 2^\circ\text{C}$  for 7 days.

### Statistical analysis

The distribution of the fungal isolates in the three communities was compared using ANOVA. P values lower than 0.05 ( $P < 0.05$ ) were considered to reflect significant differences.

### Results

The total number of four dark fungal species were isolated and identified from the sampled communities and their microscopic features are shown in figure 1. *Exophiala jeanselmei* was isolated most frequently 8 (40%), followed by *Cladophialophora carrionii* 6 (30%), *Ochroconis mirabilis* 4 (20%) while the least was *Scedosporium apiospermum* 2 (10%) (Table 1). In table 2, most of the isolates were isolated from loamy soil 13 (65%), followed by clay soil 4 (20%) while the least was sandy soil 3 (15%). Table 3 revealed that samples collected from Umuoma produced most of the dark fungi 10 (50%), followed by Umuaku 6 (30%) while the least was Aluoha 4 (20%). Statistically, the distribution of the isolates in the three communities was significant ( $P < 0.05$ ). The result of pathogenicity test revealed that all the Wistar rats infected using the fungal isolates survived, though erythematous lesions were seen at the site of infection. The scrapings collected from the infected skin yielded colonies of the inoculated fungi after incubation on SDA at  $30 \pm 2^\circ\text{C}$  for 7 days, which confirmed that the organisms were responsible for the lesions (Figure 2).



**Figure 1:** Microscopic features of dematiaceous fungi in slide culture and lactophenol cotton blue staining.

Fungal isolate	Frequency	% occurrence
<i>Exophiala jeanselmei</i>	8	40
<i>Cladophialophora carrionii</i>	6	30
<i>Ochroconis mirabilis</i>	4	20
<i>Scedosporium apiospermum</i>	2	10
Total	20	100

**Table 1:** Frequency occurrence of fungal isolates.

Fungal isolate	Loamy soil	Clay soil	Sandy soil
<i>Exophiala jeanselmei</i>	5	2	1
<i>Cladophialophora carrionii</i>	3	2	1
<i>Ochroconis mirabilis</i>	3	0	1
<i>Scedosporium apiospermum</i>	2	0	0
Total	13 (65%)	4 (20%)	3 (15%)

**Table 2:** Distribution of the fungal isolates in different soil types.

Fungal isolate	Aluoha	Umuaku	Umuoma
<i>Exophiala jeanselmei</i>	2	3	6
<i>Cladophialophora carrionii</i>	1	1	2
<i>Ochroconis mirabilis</i>	0	2	1
<i>Scedosporium apiospermum</i>	1	0	1
Total	4 (20%)	6 (30%)	10 (50%)

**Table 3:** Distribution of the fungal isolates in the three communities.



**Figure 2:** Erythematous lesions on the infected Wistar rats.

## Discussion

The need to discover sources of pathogenic fungi has become paramount in clinical mycology due to high rate of fungal infections in the society. This research has revealed that dark fungi are found in different soil types. This observation agrees with the finding of

several researchers [1-3]. The highest number of dark fungi isolated from loamy soil could be ascribed to high humus which emanated from high biodegradation as the organisms are well-known saprophytes. This conforms to the observation made by Chukwuma, *et al.* [3] who investigated the distribution and pathogenicity of dematiaceous fungi and obtained highest number of dark fungi from loamy soil. The presence of the four isolates at Umuoma community could be ascribed to variation in the habitat of the fungi. The fungi were able to adapt to environmental conditions that scared other fungal species. Similar conclusion was drawn by Chukwuma, *et al.* [3]. The ability of the isolates to produce lesions on the Wistar rats portrayed their pathogenicity potential. This confirms the previous literatures that documented that dark fungi are responsible for subcutaneous infections [3,5,6]. However, there was variation in the fungal isolates which could be ascribed to regional differences. Chukwuma, *et al.* [3] isolated from samples obtained from Awka North while samples in this study were obtained from Anambra South [7].

## Conclusion

This study has shown that dark fungi are mostly found in loamy soil due to high humus. Also, soil samples collected from Umuoma community yielded the highest number of dark fungi due to high availability of degradable organic materials. These dark fungi are pathogenic due to their ability to produce lesions on the skin of Wistar rats.

## Acknowledgment

The corresponding author sincerely acknowledges Professor Chinelo Ursula Umedium: Head of Department, Medical Laboratory Department, Chukwuemeka Odumegwu Ojukwu University- COOU, Igbariam Campus, for her awesome supervision during this project. Also, I acknowledge Ikechukwu Harmony Iheukwumere (PhD): Associate Professor in Medical Microbiology, Head of Department, Microbiology Department, COOU, and Bright Obidimma Uba (PhD): Senior Lecturer, Microbiology Department, COOU, for their financial support during this research work.

## Conflict of Interest

The authors hereby declare that there was no conflict of interest throughout the period of this study.

## Bibliography

1. Revankar SG and Sutton DA. "Melanized fungi in human diseases". *Clinical Microbiology Reviews* 23.4 (2010): 884-928.
2. Yew MS., *et al.* "A five year survey of dematiaceous fungi in a tropical hospital reveals potential opportunistic species". *Plos One* 9.8 (2014): e104352.
3. Chukwuma MC., *et al.* "Isolation of dematiaceous fungi from soil and their pathogenic potentials". *Journal of Biomedical Investigation* 6.1 (2008): 10-13.
4. Iheukwumere IH and Umedium CU. "Effects of *Gossypium hirsutum* leaf extracts on Gram negative bacteria isolated from the cervix of females with unexplained infertility". *African Journal of Medical Science* 14.1 (2013): 3261-3270.
5. Sokovic M., *et al.* "Antifungal activity of essential oils and components *in vitro* and *in vivo* on experimentally induced dermatomycoses at rats". *Digest Journal of Nanomaterials and Biostructures* 7.3 (2012): 959-966.
6. Abdel-Aziz M. "Synergism between clotrimazole and cinnamon oil: An effective (weapon) drug *in vitro* and *in vivo* against some multi drug resistant dermatophytes". *Egyptian Academic Journal of Biological Sciences G Microbiology* 4.1 (2012): 69-83.
7. Padhan DK and Pattnaik S. "*In vivo* antifungal activity of Acmella essential oil on dermatomycotic strain Trichophyton mentagrophytes (MTCC-7687)". *Der Pharmacia Sinica* 5.1 (2014): 40-44.

**Volume 22 Issue 2 February 2026**

**©All rights reserved by Osuji Malachy Ikeokwu., *et al.***