







Therefore, the sample size obtained by using the single population formula (634) is higher than the sample size calculated by using the second objective (using factors significantly associated with the outcome variable). Therefore, the minimum sample size to represent the study participants was 634.

### **Sampling technique**

Two-stage sampling techniques were used to select the study participants. First, 10 kebeles (nearly 50% of the total of 21 kebeles in Maksegnit District) were selected using a simple random sampling technique (the lottery method). The sample size was proportionally allocated to each selected kebele based on the size of the population in the kebele. Household mothers were selected by a systematic random sampling technique. The sampling interval  $K^{\text{th}}$  value was calculated by dividing the total number of mothers in each kebele by the sample size allocation.

### **Data collection tool and procedure**

The questionnaire was prepared by reviewing previous studies [25-27]. First, the English version was prepared, translated to Amharic, and then translated back to English by language experts. The questionnaire consists of four parts: socio-demographic, environmental health, knowledge toward hygiene and attitude toward hygiene. Five BSc degree holders in environmental health and one master degree holder in environmental health were recruited for the data collection process and as supervisors, respectively. Data were collected through an interview and direct observation using validated and pretested structured questionnaires.

### **Data quality management**

A pre-test was done on 5 percent of the sample size (32 participants) in Koladiba District. The questions required modifications based on the pretest. Training was given to the data collectors before the actual data collection. The training covered the aim of the study, procedures, inclusion and exclusion criteria, data collection technique (going through the questionnaires one by one), the art of interviewing, a way of collecting the data, and clarification. The supervisor was checking on how the data collection was going on a daily basis. At the end of each day of data collection, the principal investigator also checked the completeness of each questionnaire. Every questionnaire was checked before data entry by the principal investigators. Study participants were clearly informed about the purpose and usefulness of the survey, thereby creating a friendly atmosphere to reduce their stress as the study touches on sensitive issues. The validity of the questionnaire, which is measured by its type of content validity, was acquired by the experts, and its reliability was established using the internal stability method (Cronbach's alpha coefficient). A multi-collinearity assumption was checked using the Variance Inflation Factor (VIF < 1.5). Model fitness was checked using the Hosmer-Lemeshow test which for knowledge was ( $p = 0.422$ ).

### **Data processing and analysis**

Data were validated, coded, and entered into Epi-data 4.6 before being exported to SPSS 23 for cleaning and analysis. Descriptive statistics were carried out and presented with narration and tabulation. Bivariate logistic regression analyses were carried out to identify independent predictor variables for the outcome variables. Multivariable logistic regression was used to generate the final fitted mode in order to identify variables that were significantly associated with knowledge of house and environment hygiene among households. Bivariable p-values of less than or equal to 0.25 were considered to be select candidate variables for multivariable logistic regression analysis. The odds ratio with a 95% confidence interval was used to determine the strength of association between the predictors and outcome variables. A p-value of 0.05 was considered adequate to declare statistical significance.

### **Variables and measurements**

Knowledge towards house and environment hygiene was a dependent variable, while socio-demographic factors (age, ethnicity, religion, residence, marital status, and education status, occupation, family size, and access to media) and household and environmental

factors (availability of water, source of water, daily water consumption, availability of latrines, hand-washing facilities, household cleanliness, number of rooms, drinking water treatment, disposal of children’s feces, cleaning of storage containers, community health services) and attitude were independent variables. Knowledge about of hygiene respondents was asked 12 knowledge questions (Cronbach’s alpha 0.85) about their knowledge of hygiene of household mother about house and environment.

For each question; the participants were given 1 point for the correct answer and a zero (0) when the answers were incorrect. The responses were added, and the mean was computed. Those study participants who scored mean and above mean of the sum of the knowledge questions were considered as having good knowledge and otherwise they were labeled as having “poor knowledge” [22].

Attitude about house and environmental hygiene among household mothers were assessed by using 16 attitudinal questions (Cronbach’s alpha 0.72) with a 5-scale Likert (1-strongly agree, 2-agree, 3-undecided/not applicable, 4-disagree and 5-strongly disagree dichotomized as 1 for strongly agree, agree and 0 for undecided/ not applicable, disagree and strongly disagree. Those study participants who scored mean and above mean of the attitude questions were considered as holding the as favorable attitude (> mean) or unfavorable attitude (< mean) [28].

**Ethical approval and consent to participate:** An approval letter was given from Gondar University’s research and ethics committee (Ref. IPH/2274/2014). In addition, after explaining the importance of the study, permission letters were obtained from each of the kebeles/ ketenas administrators.

**Informed consent statement:** Verbal informed consent has been obtained from the patient(s) to publish this paper”.

## Results

### Socio-demographic characteristics of participants

A total of 634 mothers participated in this study, with a response rate of 100%. The mean age of the study participants was 36.7 ( $\pm$  9.9 SD). About 582 (91.8%) of the study participants were from the Amhara ethnic group. The majority of study participants, 393 (62%), were currently residing in urban areas. However, regarding the educational status of the study participants, only 63 (9.9%) of them attended secondary school or above. Furthermore, among the 634 total study participants, more than half (56.6%) do not get access to information about housing and environmental hygiene (Table 2).

Variables	Frequency	Percent (%)
<b>Age groups</b>		
18-26	87	13.7
27-35	234	36.9
36-44	185	29.2
45-70	128	20.2
<b>Ethnicity</b>		
Amhara	582	91.2
Tigray	52	8.2
<b>Religion</b>		
Orthodoxs	573	90.4
Muslim	56	8.8
Other	5	0.8

<b>Mothers' marital status</b>		
Current married	479	75.6
Single	155	24.4
<b>Husbands' educational (n = 479)</b>		
Unable to read write	171	35.7
Able to read write	230	48.0
Primary school	43	9.0
Secondary and above	35	7.3
<b>Mothers' educational</b>		
Unable to read write	387	61.0
Able to read write	145	22.9
Primary school	39	6.2
Secondary and above	63	9.9
<b>Mothers' occupation</b>		
Housewife	454	72.1
Civil servant	145	22.9
Merchant	32	5.0
<b>Family size in number</b>		
≤ 4	329	51.9
> 4	305	48.0
<b>Place of residence</b>		
Urban	393	62.0
Rural	241	38.0
<b>Access to media</b>		
Yes	378	56.6
No	256	40.4

**Table 2:** The socio-demographic factors with respect to knowledge of housing and environment hygiene among mothers in Maksegnit District, Ethiopia, 2022.  
Other: Catholic and protestant.

### Environmental related factors

Of the total 634 mothers, 368 (58%) had access to private latrines. Of these, only 119 (18.8%) mothers had latrines with hand washing facilities. Of this, only 67 (10.6%) functional hand washing facilities had water and soap during the data collection. Regarding access to a safe water supply, the majority of study participants-349 (55.0%) responds to get water from piped water sources. Furthermore, a higher proportion of mothers more than two-thirds of 464 (73.2%) got to the water within 30 minutes. Individual daily water consumption was 20 L/C/D for 522 (82.3%) of the study participants. In terms of water treatment, the majority of study participants (461, or 72.7%) did not treat water. In terms of waste disposal, nearly half of the study participants (50.5%) disposed of their waste in an open field (Table 3).

Variables	Knowledge		Frequency
	Good	Poor	
<b>Presence of latrine</b>			
Yes	275	93	368 (58.0)
No	184	82	266 (42.0)
<b>Latrine present within 6 meters in household</b>			
Yes	213	72	285 (77.4)
No	62	21	83 (22.6)
<b>Hand wash facility near to latrine</b>			
Yes	86	33	119 (18.8)
No	373	142	515 (81.2)
<b>Water source</b>			
Piped water	266	83	349 (55.0)
Protected well	47	26	73 (11.5)
Protected spring	116	46	162 (25.6)
Un protected well	7	4	11 (1.7)
Un protected spring	23	16	39 (6.2)
<b>Long to take water</b>			
≤ 30 minute	230	234	464 (73.2)
> 30 minute	129	41	170 (26.8)
<b>Water treatment</b>			
Yes	129	53	173 (27.3)
No	339	122	461 (72.7)
<b>Water treatment option</b>			
Chlorine /wuhaagar	123	52	175 (27.6)
Boiling	20	10	30 (4.7)
No usage treatment	316	113	429 (67.7)
<b>Frequency of cleaning water storage</b>			
Daily	66	26	92 (14.5)
Weekly	186	76	262 (41.3)
Monthly	47	14	61 (9.6)
When dirty	160	59	219 (34.5)
<b>Reliability of water in a year</b>			
Yes	235	80	315 (49.7)
No	224	95	319 (50.3)
<b>Amount of water used</b>			
≥20 L/C/D	28	84	112 (17.7)
<20L/C/D	234	288	522 (82.3)
<b>Waste disposal management</b>			

Open field	221	99	320 (50.5)
In the pit	161	52	213 (33.6)
Burning near the yard	77	24	101 (15.9)
<b>Have live stock</b>			
Yes	292	128	420 (66.4)
No	167	47	214 (34.6)
<b>Observed flies in the house</b>			
Yes	230	111	341 (53.8)
No	229	64	293 (46.2)
<b>Room of cooks food</b>			
Separate kitchen	136	61	197 (31.1)
Not separated kitchen	323	114	437 (68.9)
<b>Hand wash after toilet</b>			
Yes regularly /always	229	73	300 (47.6)
Yes, sometimes	196	78	274 (43.2)
Never	34	24	58 (9.1)
<b>Contact community health services</b>			
Yes	173	74	247 (39.0)
No	286	101	387 (61.0)
<b>Attitude</b>	301	91	392 (61.8)
Feverable	158	84	242 (38.2)
Unfeverable			

**Table 3:** The housing and environmental related factors with respect to level of knowledge hygiene among mothers in Maksegnit District, Ethiopia, 2022.

### Knowledge about house and environmental hygiene

After adding the responses and computed mean, the level of knowledge about house and environmental hygiene among household mothers was 72.34% CI (68.9 - 75.7).

### Factors associated with participant's level of knowledge

Socio-demographic variables included maternal age group, ethnicity, marital status, educational level of mothers, family size, residence, and access to media in a bivariable binary logistic regression analysis; environmental variables included housing end presence of latrines, type of water source, reliability of water in a year, waste disposal management practice, presence of livestock, fly presence in the house, place of cooking, and presence of animal faces near the house. From these variables, residence and access to media, fly presence in the house, and room of cooking were significantly associated with hygiene knowledge among household mothers.

In this study, urban dwellers were 1.61 times (AOR: 1.61; 95% CI [1.04, 2.49]) more likely than rural dwellers to have good knowledge about housing and environmental hygiene. Participants in the study who had access to the media were 1.82 (AOR: 1.82; 95% CI [1.17,



2.84]) times more likely to have good knowledge of housing and environmental hygiene than those who did not have access to housing and environmental hygiene information in the media. When compared to study participants whose houses had flies, those who had flies were 1.77 (AOR: 1.77; 95% CI [1.08, 2.91]) times more likely to have good knowledge about housing and environmental hygiene. As well, those study participants who make food in a separate room were 1.89 times more likely to have good knowledge toward housing and environment hygiene than those who make food inside the main living room (AOR: 1.89; 95% CI [1.13, 3.18]), and those study participants who make food in an open space outside the household were also 4.72 times more likely to have good knowledge toward housing and environment hygiene compared with those study participants who make food inside the main living room (AOR: 4.72; 95% CI [1.66, 13.56]). Finally, the probability of good knowledge was 2.94 times higher (AOR: 2.69; 95% CI: [1.33-2.82]) among participants who had a favorable level of attitude towards housing and environment hygiene (Table 4).

Variables	Knowledge level		COR (95% CI)	AOR (95% CI)
	Good	Poor		
<b>Ethnicity</b>				
Amhara	426	156	1.57 (0.87, 2.85)	-
Tigray	33	19	1	
<b>Age groups</b>				
18 - 26	69	18	1.62 (0.85, 3.08)	-
27 - 35	169	65	1.10 (0.68, 1.77)	-
36 - 44	131	54	1.03 (0.63,1.68)	-
45 - 70	90	38	1	
<b>Maternal education</b>				
Secondary and above	52	11	1.94 (0.99, 3.84)	-
Primary school	30	9	1.37 (0.64, 2.95)	-
No formal education	377	155	1	
<b>Mothers' marital status</b>				
Single	120	35	1.42 (0.93, 2.17)	-
Current married	339	140	1	
<b>Family size in number</b>				
≤ 4	249	80	1.41 (0.99, 1.20)	-
> 4	210	95	1	
<b>Residence</b>				
Urban	306	87	2.02 (1.42-2.88)	1.61 (1.04, 2.49) *
Rural	153	88	1	1
<b>Access to media</b>				
Yes	284	94	1.40 (0.98, 1.99)	1.82 (1.17, 2.84) *
No	175	81	1	1
<b>Presence of latrine</b>				
Yes	275	93	1.32 (0.93, 1.87)	-
No	184	82	1	
<b>Water source</b>				

Protected	429	155	1.85 (1.02-3.35)	-
Unprotected	30	20	1	
<b>Waste disposal management</b>				
Open field	221	99	1	
In the pit	161	52	1.39 (0.94, 2.06)	-
Burning near the yard	77	24	1.44 (0.86, 2.41)	-
<b>Have live stock</b>				
Yes	292	128	0.64 (0.06, 0.89)	-
No	167	47	1	
<b>Animal faces near the HH</b>				-
Yes	222	108	1	
No	70	20	1.70 (0.99, 2.94)	-
<b>Observed flies in the house</b>				
Yes	230	111	1	1
No	229	64	1.73 (1.21, 2.470)	1.77 (1.08, 2.91) *
<b>Room of cooks food</b>				
Inside the main living room	85	54	1	1
Separate room	323	114	1.80 (1.20,2.69)	1.89 (1.13, 3.18) *
Open space outside HH	51	7	4.63 (1.96, 10.98)	4.72 (1.66, 13.56) *
<b>Attitude</b>				
Favorable	301	91	1.76 (1.24-2.51)	2.94 (1.33-2.82) **
Unfavorable	158	84	1	1

**Table 4:** Bivariate and multivariable analysis of factors associated with knowledge of housing and environment hygiene among mothers in Maksegnit district, Ethiopia, 2022.

Key: \*= p-value < 0.05, \*\* p-value <0001, The Hosmer and Lemeshow Test P-Value was 0.422.

## Discussion

The aim of this study was to assess the level of knowledge and associated factors regarding house and environmental hygiene among mothers living in the Maksegnit District. As a result, the overall level of knowledge about house and environmental hygiene was 72.34% CI (68.9 - 75.7). This level of knowledge about house and environmental hygiene was in line with the percentages reported in literature, such as in urban areas in Northwest Ethiopia (75.7%) [29] and a study among mothers of under-five children in rural households, in Nepal (74.28%) [30]. Possible explanations for these similarities include the study participants' similar rural socio-demographic characteristics. However, the reported percentage of knowledge was higher than the previous studies in Northern Ethiopia 42.2% [22], Southern Ethiopia (62.1%) [28] and in Angolela, Ethiopia 52% [31]. These discrepancies might be due to differences in the study periods and participants. Study from in Angelia Ethiopia [31] the study participants were school-age children who are less knowledgeable about hygiene than adult study participants.

In this study, being urban, having access to media, having an absence of flies in the household, and cooking on an open space outside the main house were significantly associated with having good knowledge about hygiene among household mothers in Maksegnit District. Mothers who live in cities have a higher likelihood of having good hygiene knowledge than mothers who live in rural areas. This might be due to the fact that living in an urban area may create suitable conditions for ease of information sharing and communication due to improved access to mass media. This finding was supported by a study conducted in Tanzania [32] that discovered mothers who had access to mass media had a higher likelihood of having good hygiene knowledge than their counterparts. The possible explanation might be that media access inevitably increases their awareness of hygiene knowledge.

Similarly, those household mothers whose house was free from flies were associated with higher odds of good maternal knowledge about hygiene than mothers whose house was found to have flies. The possible reason might be that increased knowledge helps the mothers keep their home and its immediate environment clean and avoid breeding house flies. Moreover, those mothers who cooked in the open space outside the main living area had greater hygiene knowledge than those who cooked in the main house. This might be due to the fact that cooking in an open space other than the main living area might promote the hygiene and cleanliness of the living area, which are largely dependent on the presence of a separate kitchen for cooking. Finally, we also found a significant association of participants' attitudes and levels of knowledge. Another study supported this [28]. Persons with a positive attitude about house and environmental hygiene as a result of having a favorable attitude towards house and environment hygiene can often handle waste from home and the nearby.

### **Limitations of the Study**

This study was limited by its cross-sectional design, which prevented causality determination. Another limitation was that we did not include qualitative data. Thus, there might be a bias toward social desirability. Also, the tool we used in this study was not a standard.

### **Conclusion and Recommendation**

Overall, the level of knowledge was relatively low. Household mothers' good hygiene knowledge was significantly associated with urban living, access to media presence in the house, preparing food in the main living room, preparing food in open space outside the household and a favorable attitude. As a result, interventions that would address the above mentioned factors need to be implemented. The district health office should develop a health education program for both urban and rural mothers on the importance of latrines, hygiene facilities near latrines, a separate area for food preparation, access to water, and access to media on how to keep housing and the environment clean.

The low level of good knowledge can be improved by providing training for mothers, and women's health development armies' weekly discussion agendas should include the plan to raise issues related to hygienic housing and compounds to facilitate the sharing of information and increase the awareness of mothers. Therefore, health extension workers should design training programs on keeping houses and environments hygienic for strengthening participant's level of knowledge. Further research into the adequacy of sanitary services and hygiene information communication is advised.

### **Funding Support**

This research received no external funding.

### **Acknowledgments**

To begin, the authors would like to express their gratitude to the Gondar University College for approving the ethical clearance. Secondly, we would like to acknowledge the study participants, data collectors, and supervisors who participated in the study. Lastly, we want to thank our colleagues and friends for their support and constructive comments.

### **Author Contributions**

Conceptualization: Agerie Mengistie.

Data curation: Agerie Mengistie.

Formal analysis: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Funding acquisition: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Investigation: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Methodology: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Project administration: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Resources: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Software: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Supervision: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Validation: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Visualization: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Writing - original draft: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

Writing - review and editing: Agerie Mengistie, Worku Chekole, and Zemene Habtu.

### **Availability of Data and Materials**

The data sets used and/or analyzed during this study are available from the corresponding author on reasonable request.

### **Competing Interests**

The authors have as declared that they have no competing interests in any aspect of the article.

### **Bibliography**

1. Organization WH. Who water, sanitation and hygiene strategy 2018-2025. World Health Organization (2018).
2. Alasow MA., *et al.* "Knowledge, Attitudes and Practices on Sanitation Among Women of Reproductive Age at Badbado Camp, Dharkenley District, Mogadishu-Somalia". *Central African Journal of Public Health* 6.4 (2020): 220.
3. Mahmudiono T., *et al.* "Nutrition Education on Food Hygiene and Sanitation to increase Knowledge, Attitude and Practice among Canteen Food Handler in Indonesia". *Systematic Reviews in Pharmacy* 11.11 (2020): 1396-1400.
4. Yadav K., *et al.* "Knowledge, Attitude and Practice on Hygiene and Sanitation among population of selected districts in Province No. 2, Nepal". *South East Asia Journal of Medical Sciences* (2020): 1-9.
5. Nikwigize MGUaS. "Factors Associated with Hygiene and Sanitation Practices among Adults in Muhanga District, Rwanda". *EC Nursing and Healthcare* 4.6 (2022): 26-38.
6. CEPAL N. The 2030 Agenda and the Sustainable Development Goals: An Opportunity for Latin America and the Caribbean (2018).

---

**Citation:** Agerie Mengistie Zeleke., *et al.* "Knowledge and Associated Factors of House and Environment Hygiene among Household Mothers in Maksegnit District, Central Gondar Zone, Northwest Ethiopia: A Community-Based Cross-Sectional Study". *EC Nursing and Healthcare* 6.1 (2024): 10-23.

7. Cumming O., *et al.* "The implications of three major new trials for the effect of water, sanitation and hygiene on childhood diarrhea and stunting: a consensus statement". *BMC Medicine* 17.1 (2019): 173.
8. Ramesh A., *et al.* "Evidence on the effectiveness of water, sanitation, and hygiene (WASH) interventions on health outcomes in humanitarian crises: a systematic review". *PloS one* 10.9 (2015): e0124688.
9. Mohd R and Malik I. "Sanitation and hygiene knowledge, attitude and practices in urban setting of Bangalore: a cross-sectional study". *Journal of Community Medicine and Health Education* 7.04 (2017): 2-6.
10. JMP, WHO, UNICEF. Progress on household drinking water, sanitation and hygiene (2021).
11. Organization WH. Safer water, better health (2019).
12. WHO. Key Facts about Diarrhoeal disease (2017).
13. Cairncross S., *et al.* "Water, sanitation and hygiene for the prevention of diarrhoea". *International Journal of Epidemiology* 39.1 (2010): I193-i205.
14. UNICEF. Committing to child survival. A promise renewed: UNICEF progress report New York (2015).
15. Yallew WW., *et al.* "Assessment of water, sanitation, and hygiene practice and associated factors among people living with HIV/AIDS home based care services in Gondar city, Ethiopia". *BMC Public Health* 12.1 (2012): 1-10.
16. Akter T and Ali AM. "Factors influencing knowledge and practice of hygiene in Water, Sanitation and Hygiene (WASH) programme areas of Bangladesh Rural Advancement Committee". *Rural and Remote Health* 14.3 (2014): 31-40.
17. Johnson RC., *et al.* "Assessment of water, sanitation, and hygiene practices and associated factors in a Buruli ulcer endemic district in Benin (West Africa)". *BMC Public Health* 15.1 (2015): 801.
18. Tadele Yohannes AW and Henok Asefa. "A Cross Sectional Study: Availability of Improved Sanitation Facilities and Associated Factors among Rural Communities in Lemo Woreda, Hadiya Zone, Southern Ethiopia". *Open Access Library Journal* (2014).
19. Adem GDGaDB. "Knowledge, Attitude, and Practice on Hygiene and Morbidity Status among Tertiary Students: The Case of Kotebe Metropolitan University, Addis Ababa, Ethiopia". *Hindawi Journal of Environmental and Public Health* 2094621 (2018): 9.
20. Assefa Y., *et al.* "Community health extension program of Ethiopia, 2003–2018: successes and challenges toward universal coverage for primary healthcare services". *Globalization and Health* 15.1 (2019): 24.
21. Teshome HA., *et al.* "Complementary Food Feeding Hygiene Practice and Associated Factors among Mothers with Children Aged 6 and ndash;24 Months in Tegedie District, Northwest Ethiopia: Community-Based Cross-Sectional Study". *Hygiene* 2.2 (2022): 72-84.
22. Berhe AA., *et al.* "Knowledge, Attitude, and Practices on Water, Sanitation, and Hygiene among Rural Residents in Tigray Region, Northern Ethiopia". *Journal of Environmental and Public Health* (2020): 5460168.
23. Sahiledengle B., *et al.* "Sanitation practice and associated factors among slum dwellers residing in urban slums of Addis Ababa, Ethiopia: a community based cross-sectional study". *Journal of Public Health and Epidemiology* 10.10 (2018): 370-379.
24. Shehmolo M., *et al.* "Magnitude and Factors Associated with Hygiene Practice Among Primary School Children in Mareko District, Southern Ethiopia: A Cross-Sectional Study". *Journal of Multidisciplinary Healthcare* 14 (2021): 311.

25. Mamo Z Degebasa DZW and Mokonnnon T Marama. "Diarrheal status and associated factors in under five years old children in relation to implemented and unimplemented community-led total sanitation and hygiene in Yaya Gulele in 2017". *Dove Press Pediatric Health, Medicine and Therapeutics* (2018): 9.
26. Kuddus Mohammad Abdul SMAH., et al. "Study on Knowledge and Practice of WASH among Under 5 Children's Mother in Rural Community of Bangladesh". *International Journal of Rural Development, Environment and Health Research* 4.6 (2020): 2456-8678.
27. Marufa Sultana RAM., et al. "Hand hygiene knowledge and practice among university students: evidence from Private Universities of Bangladesh". *Dove Press Risk Management and Health Care Policy* (2016).
28. Afework A., et al. "Moving Up the Sanitation Ladder: A Study of the Coverage and Utilization of Improved Sanitation Facilities and Associated Factors Among Households in Southern Ethiopia". *Environmental Health Insights* 16 (2022): 11786302221080825.
29. B Abera WM., et al. "Water safety, sanitation and hygiene related knowledge, attitudes and practices among household residents in periurban areas in Northwest Ethiopia". *Ethiopian Journal of Health Development* 32.3 (2018): 1-7.
30. Rima Kumari Sah PKS., et al. "Assessment of the Knowledge, Attitude and Practice Regarding Water, Sanitation and Hygiene among Mothers of Under-five Children in Rural Households of Saptari District, Nepal". *American Journal of Public Health Research* 5.5 (2017): 163-169.
31. Alyssa Vivasa BG., et al. "Knowledge, Attitudes, and Practices (KAP) of Hygiene among School Children in Angolela, Ethiopia". *Journal of Preventive Medicine and Hygiene* 51.2 (2011): 73-79.
32. Alexander CC., et al. "Media access is associated with knowledge of optimal water, sanitation and hygiene practices in Tanzania". *International Journal of Environmental Research and Public Health* 16.11 (2019): 1963.

**Volume 6 Issue 1 January 2024**

**©All rights reserved by Agerie Mengistie Zeleke., et al.**