

Dietary Diversity, Household Food Security and Nutritional Status of Children (Aged 6 - 23 Months) in Jima Geneti District, Oromia, Ethiopia

Tamiru Yazew* and Agama Daba

Department of Food Science and Nutrition, Wollega University, Shambu, Ethiopia

*Corresponding Author: Tamiru Yazew, Department of Food Science and Nutrition, Wollega University, Shambu, Ethiopia.

Received: August 24, 2020; Published: September 29, 2020

Abstract

Introduction: The burden of under-nutrition, poor dietary diversity and food insecurity remains a global challenge in poor countries including Ethiopia, where their livelihood relies on agriculture. Therefore, this study was designed to assess dietary diversity, household food security status and nutritional status of children (aged 6 - 23 months) in Jima Geneti district, Oromia, Ethiopias.

Method: A community based cross sectional study was done in May, 2020. Multi stages sampling methods were employed to select 500 children. Child dietary diversity score (DD) was assessed using a 24 hrs recall method. Household Food Insecurity Access Scale (HFIAS) and anthropometric data were used to determine food insecurity and nutritional status, respectively. SPSS version 20.0 was used for analysis. Bivariate and multivariate logistic regression analyses were conducted to identify determinants of under-nutrition among children and statistical association was declared significant if p-value was less than 0.05.

Results: In this study, low DD and household food insecurity was 52.2% and 19.6%, respectively. The study also found that the prevalence of underweight, stunting and wasting among children was 19.6%, 27% and 11.8%, respectively. In addition, the finding of the present study showed low wealth index (AOR = 1.4, 95%CI: 2.3, 4.8) and food insecurity (AOR = 3.2, 95%CI: 2.3, 4.82) were significantly associated with underweight. Moreover, child birth order above three (AOR = 2.21, 95%CI:1.21,3.58), no ANC visit (AOR = 2.8, 95%CI:1.66,4.57), no consumption of wild edible foods (AOR = 6.8, 95%CI:3.81,14.2) and poor DD (AOR = 2.41, 95%CI:1.36,4.37) were significantly associated with stunting. Furthermore, this study indicated that having a history of sickness (AOR = 2.5, 95%CI: 2.82, 3.4), no ANC visit (AOR = 4.28, 95%CI: 2.13, 8.63) and low DD (AOR = 3.4, 95%CI: 3.1, 7.87) were factors significantly associated with wasting.

Conclusion: This study concluded that under-nutrition among children in the study area is very prevalent. Therefore, a community based nutrition sensitive agriculture interventions should be done for generating income and diversifying foods to curtail under nutrition and household food insecurity problems in the study area.

Keywords: Nutritional Status; Underweight; Stunting; Wasting; Dietary Diversity; Food Security

Abbreviations

CI: Confidence Interval; DE: Design Effect; DD: Dietary Diversity; FAO: Food and Agriculture organizations; HFI: Household Food Insecurity; HFIAS: Household Food Insecurity Access Scale ; IFPRI: International Food Policy and Research Institute; PPS: Population Proportion to Size; SRS: Simple Random Sampling Method; SSA: Sub Sahara Africa

Introduction

Adequate nutrition is essential during childhood to ensure healthy growth, proper organ development and function, a strong immune system, and cognitive development. However, the burden of under-nutrition and food insecurity remains a global challenge in sub-Saha-

ran Africa [26]. Despite the overall progress to reduce global food insecurity and chronic undernourishment, sub-Saharan Africa remains the most food-insecure region in the world with close to 223 million people undernourished [20]. This challenge could affect young children, pregnant and lactating women who are the most vulnerable and at high risk.

In Ethiopia, magnitude of under-nutrition, low dietary diversity and food insecurity is very high in rural communities where livelihood depends on farming system [43]. The nutritional consequences of food insecurity include under nutrition depending on a broad range of contextual, economic and sociocultural factors [23]. A study conducted in Tigray, Ethiopia, showed that the prevalence of stunting, underweight and wasting in food insecure households was high and 52.1%, 20.5% and 12.6%, respectively [29]. In addition, a study done in south Ethiopia [11], reported that the prevalence of stunting, underweight and wasting among children was 45.6%, 26.3%, and 14.6%, respectively. Besides, a study done in Ethiopia revealed that the prevalence of stunting, underweight and wasting was 50.7%, 27.5% and 5.9%, respectively [8]. A study done by [6] also reported that the prevalence of stunting, underweight and wasting was 42%, 33% and 15%, respectively.

Several factors which are associated with under-nutrition have also been identified; low wealth index [7,12,17,30,32,49,50]. Another factors associated to under-nutrition were food insecurity [37,45], no antenatal care visit [12] and birth order [19,41]. In addition, factor which associated with under nutrition among children was child dietary diversity score [9,28,31,36,44,51]. Besides this, diarrhea (sickness) was also another factor associated with under-nutrition [7,10,12,32,39].

Food is a primary need basic to all human needs and a fundamental human right [27]. Improved food security is vital in the alleviation of poverty, promotion of people's health and labor productivity, contributes to the political stability of a country and ensures sustainable development of citizens [20]. The Food and Agriculture Organization (FAO) defines food security as a "situation when all people at all times have physical, social and economic access to sufficient, safe and nutritious food to meet their dietary needs for an active and healthy life" [20]. However, food insecurity remains a major concern for numerous rural households in Sub-Saharan Africa and it influences children nutritional statuses by limiting the quantity and quality of dietary intake [20]. Approximately, one billion people globally are chronically hungry; two billion regularly experience periods of food insecurity [26]. This burden of under-nutrition falls disproportionately on infants and young children and women, especially pregnant and lactating, in low and middle-income countries.

According to a study done in south Ethiopia, the prevalence of household food insecurity was high (75.8%) and it was significantly associated with underweight and stunting among children under five years old [11]. Another study done in the three regions of Africa by [8], also revealed that being stunted or underweight were significantly higher for children in severely food-insecure households in Bangladesh and Ethiopia and in moderately food-insecure households in Vietnam. Moreover, a study done in west Oromia (Ethiopia), indicated that the prevalence of food insecurity was 74.1% and it had association with children underweight [37].

Dietary diversity is considered to be a key indicator in assessing the access, utilization, and quality of diet of individuals or households [20]. Since no single food can contain all nutrients, the more food groups included in the daily diet, the greater the likelihood of meeting nutrient requirements [33]. Thus, a diet that is sufficiently diverse may reflect nutrient adequacy. The diversity of species used in agricultural and livelihood systems is also essential for human nutrition and sustainable food systems [40].

Poor dietary diversity is a severe problem among poor communities, especially among children, pregnant and lactating mothers because they require additional energy and nutritious foods for their physiological and mental development [38]. Because low-quality diets are the leading risk factor for ill health worldwide [1] and determined by socioeconomic and other factors including income, education, gender empowerment, and inequality [16]. Studies done in Ethiopia revealed that almost above half percentages of children under five years had poor dietary diversity score [14,15,28,34,47,48], in Tanzania [31] and in Burkina Faso [39]. Another study done in southwest Ethiopia also shows that only 38% of children met the minimum dietary diversity score [18].

Oromia region is among the regions producing adequate food products in the Ethiopia. Nevertheless, it is reported that there is high prevalence rate of child under nutrition as compared to prevalence rates in less productive regions of Ethiopia [25]. The Ethiopian mini demographic and health survey reported that the prevalence of children stunted, underweight and wasted was 37.5%, 22.2% and 6.9%, respectively in Oromia region [13]. In rural communities, this prevalence of stunting among children in food insecure households was very high (61.1%) compared to those children who lived in food secure households [2]. However, in the study area, there was no study done to assess dietary diversity, household food security and nutritional status of children. Therefore, this study was designed to assess dietary diversity, household food security status and nutritional status of children (aged 6 - 23 months) in Jima Geneti district, Oromia, Ethiopia.

Methodology

Study area

The study was carried out in Jima Geneti district, Horo Guduru Wollega zone, Oromia region, Ethiopia. Jima Geneti is one part of the Horo Gudru Wollega Zone woredas and is bordered on the east by Guduru, on the north by Horo, on the west by east Wollega Zone, on south by Wollega Zone and on the southeast by Jimma Rare. The 2007 national census reported a total population for this woreda of 64,158, of whom 31,756 were men and 32,402 were women; 6,966 (10.86%) of its population were urban dwellers.

Study design and period

A community based cross sectional study design was conducted between May to June 2020.

Source and study population

All children aged 6 - 23 months in rural communities of Jima Geneti district, Horo Guduru wollega zone were the source population. All children aged 6 - 23 months in rural communities who were randomly selected from Jima Geneti kebeles were constituted the study population.

Inclusion and exclusion criteria

Households which are the residents of the study area at least for 6 months and which have children 6 - 23 months were included in the study area but those farmers/mothers who are chronically sick and not found at home during the time of study were excluded from the study.

Sample size determination

The sample size of the study is calculated using formula for a single population proportion by considering the following assumptions:

$$n = Z_{\alpha/2}^2 \frac{P[1-P]}{d^2}$$

Where: n = required sample sizes, $Z_{\alpha/2}$ = critical value for normal distribution at 95%CI which equals to 1.96 (z value at $\alpha = 0.05$) P = expected prevalence of stunting in Oromia region (27%) [13]. d = margin of error 5%). The formula is yielded n = 303. Therefore, the total sample size required for this study is 500 children by considering 10% of non-response rate and design effect (DE) of 1.5.

Sampling procedures

Jima Geneti district is purposively selected as study site since there was no study done before in the study area regarding assessment of dietary diversity, household food security and nutritional status of children. Therefore, multi-stages sampling methods will be used to draw samples for the study.

From the total fourteen (14) kebeles found in the Jima Geneti district, five kebeles (Gudetu Jima, Hunde Gudina, Gudetu Geneti, Lalisa Biya and Kelela Didimtu) were selected using simple random sampling technique. The children in each kebele were also selected using population proportion to size based on the available data at Horo Guduru Wollega zonal health office. Finally, from each kebele, the eligible children required for the study were selected randomly (Figure 1).




Figure 1: Schematic presentation of sampling procedure in Jima Geneti district, Ethiopia, 2020.

Data collection procedures for survey

The data about socio-demographic and economic, dietary diversity, household food security status and nutritional status in the study area were collected using a structured questionnaire adapted from different relevant studies. The questionnaire was first developed in English and translated into Afan Oromo (a survey language). Then it was translated back to English by different language experts to check for consistency. After training was given for data collectors consecutively for three days, the data were collected by ten enumerators, two supervisors and researchers. At the end of each day, the completeness of questionnaires was checked by the principal investigator.

Socio-demographic and economic data

The data about socio-demographic (sex, and age of child, parents educational and occupational status, marital status) and economic factors (wealth index, ownership of land or cattle) were collected from mothers/caregivers.

Dietary diversity data

The Diet Diversity Score and a 24-hour recall method was conducted with mothers regarding child diet intake. Mothers/caregivers are requested to list all the foods consumed by the family both at home and out of the home in the 24 hours preceding the interview. The

seven food groups were used for children aged 6 - 23 months. Considering four food groups as the minimum acceptable dietary diversity, a child with a DDS of < 4 , was classified as poor dietary diversity and high if a child DDS was ≥ 4 .

Household food (in) security data

Household food insecurity (HFI) was measured by using the Household Food Insecurity Access Scale (HFIAS) [21]. The mothers were asked nine (9) questions related to the households experience of food insecurity in the past 12 months preceding the survey. Each item starts with an occurrence question that identifies if the condition has been experienced in the household. An affirmative answer was then followed by a frequency of-occurrence question to determine if the condition happened rarely (once or twice), sometimes (3 - 10 times), or often (> 10 times) during the reference period. The responses were coded as 0 = never, 1 = rarely, 2 = sometimes, or 3 = often. From these questions, two indicators of household food insecurity (HFI) were constructed: Household Food Insecurity Access Prevalence (HFIAP) score, which ranges from 0 to 27. The HFIAP indicators were categorized households into four (4) levels of HFI: food-secure, mild, moderately, and severely food-insecure [21]. For this study, only two levels of Household Food Insecurity Access Scale (food secure and insecure) were used since the sample size was small.

Anthropometric data

The weight of the child was measured using electronic digital weight scale (to the nearest 0.1 kg) with minimum/lightly/clothing and no shoes. Calibration was done before weighing every child by setting it to zero using 5 kg of known iron metal. The height of the child was also measured without shoes (to the nearest 0.1cm) by using a vertical wooden height board. This was done by placing the child on the measuring board, and standing upright in the middle of board and then the child's head, shoulders, buttocks, knees and heels touching the board. The age of the child was calculated in months from their birth date to the day of data collection using a local event. Mothers are asked whether the child is born before or after certain major events until accurate age is pinpointed.

Variables of the study

In this study, under-nutrition (stunting, underweight and wasting) of children was used as dependent variables while socio-demographic and economic factors, household food security status, child dietary diversity score were used as independent variables.

Data analysis

Data were entered in to SPSS statistical software version 24.0. Before the analysis, data were checked for completeness and then cleaned. Descriptive statistics was run using tables, graphs, and percentage points was calculated to describe household food security status, dietary diversity score and nutritional status of children in the of study area. Height for age, weight for age and weight for height of children were generated from WHO growth standards using WHO Anthro program, version 3.2.2. A child was considered as stunted, underweight and wasted if Z score was $< - 2$ for each index. Both bivariate and multivariate logistic regression was used to identify factors associated to nutritional status of children and p-value less than 0.05 was considered as statistically significant. The degree of association between dependent and independent variables was reported using adjusted odds ratio (AOR) and 95% CI.

Results

Socio-demographic and economic characteristics of participants

A total of 500 households were included in this study with a response rate of 100%. Of the total participants, about 472 (94.4%) mothers were married. Four hundred six (81.2%) mothers had no formal education. This study also shows that about 313 (62.6%) participants had family size greater than five. in the study were Protestant and Orthodox religion, respectively. Regarding the wealth status, about

296 (59.2%), 113 (22.6%) and 91 (18.2%) households were categorized as low, medium and high wealth index, respectively. The details information of background and economic characteristics of the respondents is listed in the below.

Variables	Frequency	Percentage (%)
Age of the child (in months)		
6 - 8 months	185	37
9 - 11 months	212	42.4
12 - 23 months	103	20.6
Sex of the child		
Male	203	40.6
Female	297	59.4
Birth order of the child		
First	86	17.2
Second	99	19.8
Third and above	315	63
Maternal age (in years)		
< 25 years	124	24.6
25 - 36 years	218	43.6
≥ 36 years	158	31.6
Religion		
Orthodox	82	16.4
Muslim	117	23.4
Protestant and others	301	60.2
Marital status of the mother		
Single	8	1.6
Married	472	94.4
Divorced and others	20	4
Education status of the mother		
Attend formal education	94	18.8
Not attend formal education	406	81.2
Occupational status of the mother		
Housewife	369	73.8
Farmer	12	2.4
Gov't employment	24	4.8
Merchant	57	11.4
Daily labor	38	7.6
Education status of the mother		
Attend formal education	123	24.6
Not attend formal education	377	75.4

Occupational status of the father		
Farmer	371	74.2
Gov't employment	56	11.2
Merchant	29	5.8
Daily labor	44	8.8
Household family size		
< 5	187	37.4
≥ 5	313	62.6
Household wealth index		
Low	296	59.2
Medium	113	22.6
High	91	18.2

Table 1: Socio-demographic and economic characteristics of participants at Jima Geneti district, Ethiopia, 2020.

Child dietary diversity scores

The finding of this study reported that about 469 (93.8%) participants had no home gardens while 151 (30.2%) children consumed wild edible foods during the past 24 hrs before the survey. The below table 3 also shows that almost about 428 (85.5%) and 398 (79.6%) children consumed cereals and legumes based foods, respectively. Moreover, this result reports that about 413 (82.6%) children did not get meat/poultry/fish during the survey in the study area. Concerning the dietary diversity scores of children, about 239 (47.8%) and 261 (52.2%) children had high (≥ 4) and low (≤ 3) dietary diversity scores, respectively (Table 2).

Food groups	Frequency	Percentage (%)
Did you have home gardens?		
Yes	31	6.2
No	469	93.8
Cereals/Root/Tubers		
Yes	428	85.5
No	72	14.4
Legumes/Nuts		
Yes	398	79.6
No	102	20.4
Milk, dairy and dairy products		
Yes	273	54.6
No	227	45.4
Meat/Fish/Poultry		
Yes	87	17.4
No	413	82.6

Eggs		
Yes	223	44.6
No	277	55.4
Other fruits and vegetables		
Yes	114	22.8
No	386	77.2
Vitamin A rich fruits and vegetables		
Yes	93	18.6
No	407	81.4
Wild edible foods		
Yes	151	30.2
No	349	69.8
Dietary diversity scores		
High (≥ 4)	239	47.8
Low (≤ 3)	261	52.2

Table 2: Dietary diversity Scores of the study children at Jima Geneti district, Ethiopia, 2020.

Household food security status

Based on the responses to the nine generic occurrence questions, about 143 (28.6%) households were unable to eat the kinds of foods he/she preferred during the past thirty days. This study also reports that about three hundred thirteen (62.6%) and one hundred twenty three (24.6%) households ate a smaller meal than he/she felt was needed and ate fewer meals in a day, respectively. In addition, the below table 4 also shows that about 106 (21.2%) households went to sleep at night hungry. Regarding, household food security status, about 402 (80.4%) vs 98 (19.6%) households were household food secure and insecure, respectively (Table 3).

Variables	Frequency	Percentage (%)
Worried about not having enough food		
Yes	86	17.2
No	414	82.8
Not able to eat the kinds of foods he/she preferred		
Yes	143	28.6
No	357	71.4
Ate just a few kinds of food day after day		
Yes	119	23.8
No	381	76.2
Ate food that he/she preferred not to eat		
Yes	247	49.4
No	253	50.6
Ate a smaller meal than he/she felt was needed		
Yes	313	62.6
No	187	37.4

Ate fewer meals in a day		
Yes	123	24.6
No	377	75.4
No food at all		
Yes	43	8.6
No	457	91.4
Went to sleep at night hungry		
Yes	106	21.2
No	394	78.8
Household Food security status		
Food secure	402	80.4
Food insecure	98	19.6

Table 3: Household food security status of the participants in Jima Geneti district, Ethiopia, 2020.

Nutritional status of children

The below figure 2 shows that the weight for age z-score, height for age z-score and weight for height z-score among children aged 6 - 23 months was 19.6%, 27% and 11.8%, respectively.




Figure 2: Nutritional status of children in Jima Geneti district, Ethiopia, 2020.

Factors associated to child nutritional status

In this study, wealth index and food security status were factors associated with underweight (Table 4). While birth order and wild edible food intake were significantly associated with stunting (Table 5), child sickness was associated with wasting in this study (Table 6). However, both child dietary diversity score and antenatal care visit were associated with stunting and wasting (Table 5 and 6).

Variables	Stunting Status		COR	AOR
	Stunting, N (%)	Normal, N (%)	(95%CI)	(95%CI)
Family size				
< 5	153 (56.3)	122 (53.5)	1	1
≥ 5	119 (43.7)	106 (46.5)	0.61 (0.47,1.76)	1.25 (0.43,2.54)
Child sickness in the past two weeks				
Yes	178 (61.2)	97 (46.4)	1	1
No	113 (38.8)	112 (53.6)	2.21 (1.48,3.86)*	2.5 (0.82,3.4)
Wealth index				
High	40 (16.6)	135 (52.1)	1	1
Medium	91 (37.7)	48 (18.5)	1.14 (0.25, 3.81)	1.59 (0.91,3.23)
Low	110 (45.6)	76 (29.3)	1.83 (1.14,3.27)*	1.61 (0.93,3.83)
Birth order of the child				
First	82 (35.6)	92 (34.1)	1	1
Second	33 (14.3)	72 (26.7)	0.82 (0.61,1.46)	1.31 (0.71,2.16)
Third and above	115 (0.5)	106 (39.3)	3.73 (2.78,4.01)*	2.21 (1.21,3.58)**
ANC visit				
Yes	200 (67.1)	68 (33.7)	1	1
No	98 (32.9)	134 (66.3)	2.32 (1.60,3.56)*	2.8 (1.66,4.57)**
Wild edible food consumption				
Yes	107 (46.0)	113 (42.3)	1	1
No	126 (54.0)	154 (57.7)	5.75 (4.48,9.43)*	6.8 (3.81,14.12)**
Dietary diversity scores				
Good	234 (72.9)	71 (39.66)	1	1
Poor	87 (27.1)	108 (60.33)	1.73 (1.12,2.64)*	2.41 (1.36,4.37)**
Food security				
Food secure	62 (24.7)	163 (65.5)	1	1
Food insecure	189 (75.3)	86 (34.5)	2.3 (1.21,3.71)*	0.75 (0.38,1.85)

Table 5: Factors associated with stunting among children in Jima Geneti District, Ethiopia, 2020 (n=500).

* P- value <.2 in the bivariate analysis, ** P-value< .05 in the multivariate analysis, 1= References, COR: Crude Odds Ratio; AOR: Adjusted Odds Ratio.

Discussion

Child malnutrition continues to be a major public health problem in developing countries including Ethiopia. Therefore, this study provided the opportunity to assess dietary diversity, household food security and nutritional status of children aged 6 - 23 months in Jima Geneti District, Horo Guduru Wollega Zone, Ethiopia.

Variables	Wasting Status		COR	AOR
	Wasting, N (%)	Normal, N (%)	(95%CI)	(95%CI)
Family size				
< 5	84 (36.7)	164 (60.5)	1	1
≥ 5	145 (63.3)	107 (39.5)	2.76 (1.28, 6.79)*	2.32 (0.95, 4.62)
Child sickness in the past two weeks				
Yes	156 (58.4)	95 (40.8)	1	1
No	111 (41.6)	138 (59.2)	2.4 (1.58,4.66)*	2.5 (2.82,3.4)*
Timely started complementary feeding				
Yes	153 (61.4)	119 (47.4)	1	1
No	96 (38.5)	132 (52.6)	3.24 (2.35, 6.81)*	2.9 (0.91,7.25)
Birth order of the child				
First	62 (31.0)	102 (34.0) 300	1	1
Second	30 (15.0)	92 (30.7)	0.72 (0.41,4.96)	2.31 (0.94,3.26)
Third and above	108 (54.0)	106 (35.3)	2.43 (1.78,3.31)*	2.21 (0.98,3.88)
ANC visit				
Yes	215 (69.8)	70 (36.5)	1	1
No	93 (30.2)	122 (63.5)	4.96 (3.49, 9.56)*	4.28 (2.13, 8.63)**
Wild edible food consumption				
Yes	180 (65.2)	77 (34.4)	1	1
No	96 (34.8)	147 (65.6)	2.69 (1.51, 4.79)*	1.2 (0.83, 3.82)
Dietary diversity scores				
Good	82 (57.3)	114 (32.0)	1	1
Poor	61 (42.6)	243 (68.1)	4.43 (1.42,9.94)*	3.4 (3.1,7.87)**
Food security				
Food secure	31 (13.4)	182 (68.0)	1	
Food insecure	201 (86.6)	86 (32.0)	2.27 (0.93, 5.55)	-

Table 6: Factors associated with wasting among children in Jima Geneti District, Ethiopia, 2020 (n=500).

* P- value <.2 in the bivariate analysis, ** P-value< .05 in the multivariate analysis, 1= References,

COR; crude odds ratio, AOR; adjusted odds ratio

The current study illustrated that about 19.6% of households in the study area was food insecure. The prevalence of this setting was lower than studies done in the three regions of Africa [8] and Ethiopia [4,11,22,37,42,52]. However, this study result was higher compared to a study done in New Zealand [35]. This situation could be due to the variation in sample size of the survey, ecological and economic status of the surveyed communities.

Considering dietary diversity (DD) score of children, the current study revealed that almost above half (52.2%) of children in this study had low dietary diversity. This finding was lower compared to other studies done in Ethiopia [14,15,28,34,47,48], in Tanzania [31] and in Burkina Faso [39]. However, the current prevalence of this study was higher than studies done in Ethiopia [18] and India [5,46].

According to the finding of this study, the prevalence of wasting, stunting and underweight was 11.8, 27.1% and 19.6%, respectively. This finding was found to be higher compared to studies done in Vietnam [8] and Ethiopia [24,52]. However, it was lower than studies done in Bangladesh [8] and Ethiopia [2,3,11,29,36,47]. This could be due to the present survey was conducted in the postharvest season which may increase food availability, accessibility and utilization by communities at household level and further lower the prevalence of under-nutrition.

In this study, children who lived in households of low wealth index were 1.4 times more likely to underweight compared to those who lived in high wealth index (AOR = 1.4, (95%CI: 2.3, 4.8)). The result of this study was in line with studies done in Ethiopia [49]. The finding of the present study was also agreed with other studies done in Ethiopia [12,19,29], in Bangladesh [17,32] and Sub-Sahara Africa [7]. The finding of the current study also showed that children who lived in household food insecure were 3.2 times more like to underweight than those who lived in food secure (AOR = 3.21.4, (95%CI: 2.3,4.82)). This finding was similar with studies done in in Ethiopia [11,37] and Nepal [45]. This could be due to the fact that food secure households are able to food access and provide it for their family.

Regarding stunting, this study found that children who had birth order three and above were 2.21 times more likely to stunted compared to those children who their birth order was first (AOR = 2.21, (95%CI: 1.21,3.58)). This finding was consistent with studies done in Ethiopia [19] and Bangladeshi [41]. In addition, this study implied that those children who had low DD (≤ 3) were 2.4 times more likely to stunted compared to those children who had DD ≥ 4 (AOR = 2.41, (95%CI: 1.36, 4.37)). The finding of this study was in line with studies done in in Ghana [8], Ethiopia [28,36], in Tanzania [31], in Burkina Faso [44] and in Nigeria [51], The result of this current study also found that children who did not consume wild edible food were 6.8 times more likely to stunted compared to those who did (AOR = 6.8, (95%CI: 3.81, 14.12)). This could be due to the fact that wild edible foods are rich in micronutrients and vital for growth and mental development and further it used in preventing malnutrition problem.

In the multivariate analysis of this study, children who had sickness before the survey were 2.5 times more likely to wasted compared with those who had no sickness (AOR = 2.5, (95%CI: 2.82, 3.4)). The result of this study was in line with studies done in Sub-Sahara Africa [7], in Ethiopia [10,12], in Bangladesh [32] and in Burkina Faso [39] This could be due to the sickness can lead children to weight loss and further increase the prevalence of wasting. Moreover, this present study revealed that mothers who did not follow up antenatal care had children who were 4.28 times more likely to wasted (AOR = 4.28, (95%CI: 2.13, 8.63)). The finding was in line with a study done in Ethiopia [12], which reported that mothers who did not follow antenatal care visit had children who were more likely to wasted compared those mothers who did it. Furthermore, this study implied that children who consumed low DD were 3.4 times more likely to wasted (AOR = 3.4, (95%CI: 3.1, 7.87)), compared to those who had good dietary diversity ≥ 4 food groups. This finding was similar with the study result done in northern Ghana [9].

Conclusion

This study demonstrated that under-nutrition is an important public health issue for children aged 6 - 23 months in the study area, Jima Geneti District, Ethiopia. The prevalence of childhood under-nutrition (underweight, stunting and wasting) is patterned by various factors such wealth index, food insecurity, birth order, antenatal care visit, wild edible food consumption, dietary diversity score and child sickness. Thus, nutritional programs with effective multi-sectoral approaches might be designed and should be prioritized for tackling childhood under nutrition in the study area particularly and in the zone at large level. A joint effort by the government and non-government organizations is necessary to improve the childhood nutritional status in the communities.

Conflict of Interests

The authors declare that they have no competing interests.

Ethical Consideration

Ethical clearance was obtained from Institutional Review Board (IRB) of the Wollega University, College of Health Sciences. A formal letter of permission was written to Regional Health Office, Horro Guduru Wollega Zonal Health Department, Jima Geneti Woreda Health Office, and to each selected kebele administration. After explaining the purpose of the study, verbal consent was obtained from each study participant. Participants were informed that participation is on voluntary basis. Personal identifiers were not included in the questionnaires to ensure participants' confidentiality. Nutrition education regarding dietary diversity and their sources were given by the data collectors for households with malnourished children during the survey.

Availability of Data and Materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Funding

The study was financially supported by Wollega University.

Authors' Contributions

TY conceived of the study, participated in its design, acquisition of data, analysis and interpretation of data, and drafted the manuscript and AD participated in the design of the study, acquisition of data and helped to draft the manuscript.

Acknowledgements

We thank Wollega University who funded for this research.

Bibliography

1. Abajobir A., *et al.* "Global, regional, and national comparative risk assessment of 84 behavioral, environmental and occupational, and metabolic risks or clusters of risks, 1990-2016. A systematic analysis for the Global Burden of Disease Study 2016". *Lancet* (2017).
2. Abdurahman., *et al.* "Household Food Insecurity May Predict Underweight and Wasting among Children Aged 24–59 Months". *Ecology of Food and Nutrition* 21 (2017): 18.
3. Abel., *et al.* "Prevalence of Malnutrition and Associated Factors among Under-Five Children in Pastoral Communities of Afar Regional State, Northeast Ethiopia" A Community-Based Cross-Sectional Study". *Journal of Nutrition and Metabolism* 2019 (2019): 13.
4. Agidew A and Singh KN. "Determinants of food insecurity in the rural farm households in South Wollo Zone of Ethiopia: the case of the Teleyayen sub watershed". *Agricultural and Food Economics* 6 (2018): 10.
5. Agrawal S., *et al.* "Socio-economic patterning of food consumption and dietary diversity among Indian children evidence from NFHS-4". *European Journal of Clinical Nutrition* 73 (2019): 1361-1372.
6. Ahmed A., *et al.* "Nutritional Status of Under Five Children in Ethiopia. A Systematic Review and Meta-Analysis". *Ethiopian-Journal of Health Sciences* 27 (2017): 175.
7. Akombi B., *et al.* "Stunting, Wasting and Underweight in Sub-Saharan Africa. A Systematic Review". *International Journal of Environmental Research and Public Health* 14 (2017): 863.
8. Ali D., *et al.* "Household Food Insecurity Is Associated with Higher Child Undernutrition in Bangladesh, Ethiopia, and Vietnam, but the Effect Is Not Mediated by Child Dietary Diversity". *Journal of Community and International Nutrition* 143 (2013).
9. Ali Z., *et al.* "The effect of maternal and child factors on stunting, wasting and underweight among preschool children in Northern Ghana". *BMC Nutrition* 3 (2017): 31.

10. Asfaw M., *et al.* "Prevalence of undernutrition and associated factors among children aged between six to fifty nine months in Bule Hora district, South Ethiopia". *BMC Public Health* 15 (2015): 41.
11. Bealu B., *et al.* "Household Food Insecurity and Its Association with Nutritional Status of Children 6–59 Months of Age in East Badawacho District, South Ethiopia". *Journal of Environmental and Public Health* 2017 (2017): 17.
12. Berhanu T and Tigist T. "Risk Factors Associated with Under-Five Stunting, Wasting, and Underweight Based on Ethiopian Demographic Health Survey Datasets in Tigray Region, Ethiopia". *Journal of Nutrition and Metabolism* (2019).
13. Central Statistics Agency. Ethiopian Mini Demographic and Health Survey, Addis Ababa, Ethiopia (2014).
14. Dafursa K and Gebremedhin S. "Dietary Diversity among Children Aged 6–23 Months in Aleta Wondo District, Southern Ethiopia". *Journal of Nutrition and Metabolism* (2019): 10.
15. Dangura D and Gebremedhin S. "Dietary diversity and associated factors among children 6-23 months of age in Gorche district, Southern Ethiopia", Cross-sectional study." *BMC Pediatrics* 17 (2017): 6.
16. Darmon N and Drewnowski A. "Contribution of food prices and diet cost to socioeconomic disparities in diet quality and health: A systematic review and analysis". *Nutrition Review* 73.10 (2015): 643-660.
17. Das S and Gulshan J. "Different forms of malnutrition among under five children in Bangladesh: a cross sectional study on prevalence and determinant". *BMC Nutrition* 3 (2017): 1.
18. Edris MM., *et al.* "Determinants of Dietary Diversity Score among Children Age between 6-23 Months in Bench Maji Zone, Southwest Ethiopia". *Pediatrics and Health Research* 3.3 (2018): 10.
19. Endris N., *et al.* "Prevalence of Malnutrition and Associated Factors among Children in Rural Ethiopia". *BioMed Research International* (2017): 6.
20. FAO, IFAD, WFP. The State of Food Insecurity in the World: Meeting the 2015 International Hunger Targets: Taking stock of uneven progress Italy. Rome (2015).
21. FAO. Guidelines for measuring household and individual dietary diversity. (Version 4), Rome, Italy (2008).
22. Feyisa M. "Determinants of food insecurity among rural households of South Western Ethiopia". *Journal of Development and Agricultural Economics* 10.12 (2018): 404-412.
23. Ghattas H. "Food Security and Nutrition in the context of the Global Nutrition Transition". Technical Paper. FAO, Rome (2014).
24. Haile A and Amboma T. "Children's nutritional status and its determinants in small towns, Sebeta Hawas district, Oromia, Ethiopia". *Journal of Food Science and Nutrition* 1.1 (2018): 33-34.
25. Hirvonen A and Hodinott J. "Targeting food security interventions when "everyone is poor": The case of Ethiopia's Productive Safety Net Programme". *International Food Policy Research Institute* (2014).
26. International Food Policy Research Institute. "Leveraging Agriculture for Improving Nutrition and Health; Highlights from an International Conference". Washington, DCIFPRI (2011).
27. Ingram I. "From Food Production to Food Security: Developing Interdisciplinary, Regional Level Research; Doctoral Thesis". Wageningen University, the Netherlands (2011).
28. Jisha H and Tamiru D. "Dietary Patterns and Anthropometric Status of Under-Five Children in Arba Minch Zuria, Gamo Gofa Zone: Community-Based Cross-Sectional Study". *Advances in dairy Research* 3 (2015): 138.

29. Kaysay A., *et al.* "Nutritional Status of Children aged 6-59 months from Food Secure and Food Insecure Households in Rural Communities of Saesie Tsaeda-Emba District, Tigray, North Ethiopia: Comparative Study". *Clinical Medicine Research* 4.1 (2015): 51-65.
30. Kasaye HK., *et al.* "Poor nutrition for under-five children from poor households in Ethiopia: Evidence from 2016 Demographic and Health Survey". *PLoS ONE* 14.12 (2019).
31. Khamis AG., *et al.* "The influence of dietary diversity on the nutritional status of children between 6 and 23 months of age in Tanzania". *BMC Pediatrics* 19 (2019): 518.
32. Khanam M., *et al.* "Individual, Household and Community Level Determinants of Childhood Undernutrition in Bangladesh". *Health Services Research and Managerial Epidemiology* 6 (2019): 1-12.
33. Labadarios D., *et al.* "How diverse is the diet of adult South African?" *Nutrition Journal* 10.33 (2011).
34. Misgan L L., *et al.* "Factors Associated with Dietary Diversity among Children of Agro Pastoral Households in Afar Regional State, Northeastern Ethiopia". *Academic Journal of Pediatrics and Neonatology* 5.2 (2017): 555-715.
35. MOH. Household Food Insecurity among Children in New Zealand. Wellington: Ministry of Health (2019).
36. Motbainor A., *et al.* "Stunting Is Associated with Food Diversity while Wasting with Food Insecurity among Under-five Children in East and West Gojjam Zones of Amhara Region, Ethiopia". *PLoS ONE* 10.8 (2015).
37. Mulu E and Mengistie B. "Household food insecurity and its association with nutritional status of under five children in Sekela District, Western Ethiopia: a comparative cross-sectional study". *BMC Nutrition* 3 (2017): 35.
38. Nti C. "Dietary diversity is associated with nutrient intakes and nutritional status of children in Ghana". *Asian Journal of Medical Sciences* 2.2 (2011).
39. Paré, *et al.* "Prevalence of wasting and associated factors among 6 to 23 months old children in the Sahel Region of Burkina Faso". *Pan African Medical Journal* (2019).
40. Powell B., *et al.* "Improving diets with wild and cultivated biodiversity from across the landscape". *Food Security* 7 (2015): 535-554.
41. Rahman M. "Association between order of birth and chronic malnutrition of children: a study of nationally representative Bangladeshi sample". *Cad Saude Publica* 32.2 (2016).
42. Regassa N and Stoecker J. "Household food insecurity and hunger among households in Sidama district, southern Ethiopia". *Public Health Nutrition* 15.7 (2011): 1276-1283.
43. Sarah CB., *et al.* "Targeting Food Security Interventions: The Case of Ethiopia's Productive Safety Net Program". *International Food Policy Research Institute* (2013).
44. Siéa C., *et al.* "Dietary diversity and nutritional status among children in rural Burkina Faso". *International Health* 10 (2018): 157-162.
45. Singh A., *et al.* "Household food insecurity and nutritional status of children and women in Nepal". *Food and Nutrition Bulletin* 35.1 (2014).
46. Singh SK., *et al.* "Socioeconomic Determinants of Dietary Diversity among Rural Households in Jaunpur District of Uttar Pradesh, India". *Asian Journal of Agriculture and Food Sciences* 8.3 (2020).
47. Tamiru Y and Agama D. "Feeding Practices and Nutritional Status of Children (Aged 2-5 Years) in Food Secure and Insecure Households in Kuyu Woreda, Oromia, Ethiopia". *Journal of Food and Nutrition Sciences* 8.3 (2020): 55-62.

48. Tegegne M., *et al.* "Factors associated with minimum meal frequency and dietary diversity practices among infants and young children in the predominately agrarian society of Bale zone, Southeast, Ethiopia: a community based cross sectional study". *Archives of Public Health* 75 (2017): 53.
49. Tekile A., *et al.* "Prevalence of malnutrition and associated factors among under-five children in Ethiopia: evidence from the 2016 Ethiopia Demographic and Health Survey". *BMC Research Notes* 12 (2019): 391.
50. Tosheno D T, *et al.* "Risk Factors of Underweight in Children Aged 6–59 Months in Ethiopia". *Journal of Nutrition and Metabolism* 2017 (2017): 8.
51. Ukegbu PO and Ogu VC. "Assessment of Dietary Diversity Score, Nutritional Status and Socio-demographic Characteristics of Under-5 Children in Some Rural Areas of Imo State, Nigeria". *Malaysian Journal of Nutrition* 23.3 (2017): 425- 435.
52. Wondu G. "Household Food Insecurity Predicts Childhood Undernutrition: A Cross-Sectional Study in West Oromia (Ethiopia)". *Journal of Environmental and Public Health* 2020 (2020): 9.

Volume 15 Issue 10 October 2020

© All rights reserved by Tamiru Yazew and Agama Daba.