

Updates in Shoulder Rehabilitation

Nithin Sunku*

M.B.B.S, D.Ortho, DNB Ortho, MNAMS, FAGE, FIJR, FIAA (UK & Ireland), FASM, IOA-WOC Fellow, SICOT Fellow.

Consultant Orthopaedic Joint Replacement & Arthroscopy Surgeon, Global Hospital Trauma Centre, Abu Road, Rajasthan, India

***Corresponding Author:** Nithin Sunku, M.B.B.S, D.Ortho, DNB Ortho, MNAMS, FAGE, FIJR, FIAA (UK & Ireland), FASM, IOA-WOC Fellow, SICOT Fellow, Consultant Orthopaedic Joint Replacement & Arthroscopy Surgeon, Global Hospital Trauma Centre, Abu Road, Rajasthan, India.

Received: November 15, 2023; **Published:** November 27, 2023

In rotator cuff repairs we have different techniques, among them transtendinous rotator cuff repair is being done for symptomatic partial-thickness rotator cuff tears. This is especially to have better biomechanical outcomes. In spite of that one of the hindrances we face is early postoperative stiffness and slower recovery. McBroom, *et al.* did a study of 61 patients in which accelerated physical therapy protocol (passive and active-assisted range of motion at 2 to 4 weeks, active range of motion as early as 4 to 6 weeks, strengthening at 6 to 8 weeks, and earlier discontinuation of obligatory sling wear) and found significant improvement in active range of motion at 6 weeks and 3 months postoperatively compared with standard physical therapy [1]. So accelerated PT gave good results. There were no higher rates of retear among the cohort.

In case of massive rotator cuff tears electromyographic biofeedback-guided rehabilitation is being done. A study was done by Tiryaki, *et al.* to know the effectiveness of electromyographic biofeedback-guided rehabilitation. 46 adults were divided into 2 groups one of them as control group. Then at the end of 1-year outcome measured by American Shoulder and Elbow score, shoulder flexion strength, shoulder range of motion, Numeric Pain Rating Scale, and Global Rating of Change Scale showed significant difference in the change in shoulder flexion strength and patient satisfaction from baseline. Hence concluded that electromyographic biofeedback can be used as an alternative method that increases patient satisfaction in exercise protocols [2].

Another is commonly seen primary adhesive capsulitis of the glenohumeral joint. Although it is considered a self-limiting condition recovery is not full. So, manipulation under anesthesia (MUA) can be considered. Van der Stok, *et al.* did a study comparing with diving in 3 groups MUA alone, MUA with capsular distension, and MUA with capsular distension and countertraction. At 6 months evaluated with re-MUA rate, Constant-Murley shoulder score, and visual analog scale score and found that MUA with capsular distension and countertraction reduced the need for a second MUA and resulted in a faster improvement in functional outcome and reduction of pain compared with the other 2 groups [3].

Bibliography

1. McBroom TJ, *et al.* "Accelerated versus standard physical therapy in patients with transtendinous rotator cuff repair: a propensity-matched cohort study". *Journal of Shoulder and Elbow Surgery* 31.6S (2022): S123-S130.
2. Tiryaki P, *et al.* "Effectiveness of exercises with electromyographic biofeedback in conservative treatment of massive rotator cuff tears: a randomized controlled study". *American Journal of Physical Medicine and Rehabilitation* 102.5 (2023): 419-426.

3. van der Stok J, *et al.* "Does capsular distension and a short period of countertraction improve outcome following manipulation under anesthesia for the treatment of primary adhesive capsulitis of the glenohumeral joint?" *Journal of Shoulder and Elbow Surgery* 31.4 (2022): 772-781.

Volume 14 Issue 10 December 2023

©All rights reserved by Nithin Sunku.