

Barriers to Reporting Medication Errors among Nurses: A Review Paper

Basheer Al-zubi¹, Mohammad AlBashtawy^{2*}, Mohammad Suliman³, Omar Khraisat⁴, Samer Al Haliq⁵, Ahmad Al-Bashaireh⁴ and Abdullah Alkhalwaldeh⁶

¹Master Student, Adult Health Nursing, Princess Salma Faculty of Nursing, AL Al-Bayt University, Jordan

²Professor, Princess Salma Faculty of Nursing, AL Al-Bayt University, Jordan

³Associate Professor, Princess Salma Faculty of Nursing, AL Al-Bayt University, Jordan

⁴Assistant Professor, Faculty of Nursing, Al-Ahliyya Amman University, Amman, Jordan

⁵Assistant Professor, Department of Emergency Medical Care, Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia

⁶Associate Professor, Faculty of Nursing, Jerash University, Jerash, Jordan

***Corresponding Author:** Mohammed AlBashtawy, Professor, Princess Salma Faculty of Nursing, AL Al-Bayt University, Jordan.

Received: June 01, 2020; **Published:** July 07, 2020

Abstract

Medication administration errors (MAEs) may threaten patients live, and there is no agreement about medication error definition. An electronic searching was conducted in the databases: Cochrane database for systemic review, CINAHL, Google scholar, Pub Med, and EBSCO. The review paper showed that the barriers to self-reporting of MAEs were recognized to be the fear reasons and administrative/management reasons. There is many interventions that need to be applied in hospitals to thrive low rate of medication errors and encourage nursing to report, more study is necessary to verify the effectiveness of each intervention in the prevention and reducing of medication errors rates.

Keywords: Barriers; Medication Administration Errors; Nursing; Self-Reporting

Introduction

Globally, the MAEs represent a major health problem in hospitals that cause injury and death [1] and threats patient safety [2]. Also, it cost the world about 1% of the total global health expenditure [3].

There is no agreement about the definition of medication error. Medication error is “any preventable event that may cause or lead to inappropriate medication use or patient harm while the medication is in the control of the health care professional, patient, or consumer. Such events may be related to professional practice, healthcare products, procedures, and systems, including prescribing; order communication; product labeling, packaging, and nomenclature; compounding; dispensing; distribution; administration; education; monitoring; and use” [4].

According to literature, there are different types of medication errors; with the most common type was incorrect time for administration of medication [5,6]. The other types of medication errors include: the wrong dosage calculation [7], incorrect medication [8], and incorrect patient [8], error in technique [9], omitting a drug due to drug availability [6], and administration of not prescribing drug [9]. In order to reduce MAEs and improve patients' safety, nurses should follow the six rights of medication administration [10].

There are many different factors that may contribute to MAEs; nurse-related factors (age, experience, and knowledge and skills about medications) [11], physician-related factors (such as: not clear of change medication order) [12], nurse-physician related factor (such as: poor communication between nurses and physicians) [5], organizational factors (overworked, nurse to patients ratio, night shift, and MAEs depending on the department) [13] and patients-related factors (patient age frequently in pediatric patients) [13].

The major barriers of reporting the medication errors were: Fear of negative consequences (punishment and lose of job), and fear of blame [14,15]. Other barriers that may affect the self-reporting of medication errors were: lack of policies to report, managerial/administrative factors, ambiguous process to report, nurses with high experience, level of nurses education and the nature of error [14,15].

Aim of the Study

This paper intends to review the major literature to recognize the barriers to self-reporting for MAEs. Additionally, provide recommendations to reduce the MAEs and encourage nursing to report.

Methods

Search methods

The electronic searching was conducted in different database: Cochrane database for systemic review, CINAHL, Google Scholar, Pub Med, and EBSCO. Key search terms used were: 'Barriers', 'medication administration errors', 'nursing' and 'self-reporting'.

Inclusion criteria were studies (a) discuss the medication administration errors and the barriers to report (b) published between 2007 and 2020, (c) published in English. While, review articles, comments, editorials, letters, books, and thesis reports were excluded from this study.

Search outcome

Searching of literature yielded about 40 studies for review, 20 excluded when reviewed the title, 9 when reading the abstract and 11 studies met the inclusion criteria. This final examination resulted in 10 studies, excluding studies consisting only of abstracts, review studies and irrelevant studies.

Results and Discussion

Administration of medication to patients is an important role of the nursing practice with high possibility of medication errors to occur. According to past studies there were no important differences in what nurses believed as MAEs among nurses [16] and nurses' attitudes towards MAEs reporting were tending towards positive [17]. However, the reporting among nurses was low [16]. For example, a study conducted in Ethiopia found that the prevalence of MAEs reporting was 29.1% among nurses [18]. In Taiwan, a cross-sectional study found that about 88.9% of MAEs were reported orally, while 19.0% were reported through the hospital internet system [17]. While, in Iran the prevalence of MAEs was around 17.0% [10].

Regarding reporting of medication errors, the nurses don't report errors because of the personal perceptions and misunderstanding on what constitutes an error, whether significance of the error changes the requirement that all errors should be reported [19], and the nurses didn't think that an error was serious enough [15].

The main barriers to self-reporting for MAEs found in the present review were: First, fear (fear of punishment, fear of blame, fear of annual evaluation affecting, and fear from reactions of patients and their families), [14-16,20,21] and fear related to judicial issues [17] and from disciplinary actions [15]. Second reason, administrative barriers (administrative attitudes, and response to MAEs) [14,18] and management barriers (inappropriate reaction from manager, presence of a culture of blame and disciplinary action, and inappropriate feedback of manager after error reporting) [21]. The third reason was the lack of information and difficulty to reporting errors (the

importance of error reporting, how to report, and recognize if MAEs occurs) [20,21]. Fourth, the lack of medication error recording and reporting system in hospital, lack of appropriate feedback, and lack of clear definition of MEs [21,22]. Finally, other less significant reasons associated with lack of time to reporting errors [20]; and the workload as a result of high number of patients [10].

On the other hand, some studies found an important correlation among barriers to report and a number of demographic data (gender, age, educational level, and years of experience) [23-31].

Conclusion

The purpose of the present review paper was to recognize barriers to reporting MAEs. The MAEs may be not reported or underreported because of a range of factors, including individual perceptions, fears, and misapprehension on what constitutes a medication error. Barriers to self-reporting of MAEs were recognized to be the fear reasons and administrative/management reasons. It is apparent that there are many of interventions that need to be applied in hospitals units to maintain low rate of medication errors and encourage nursing to report such errors. Because of the significance of the topic, more study is necessary to verify the effectiveness of each measure in the prevention and reducing of medication errors rates.

Implications and Recommendations for Nursing Management

- Nursing managers address the heavy workload to reduce MAEs.
- Nursing managers work in providing updated medication administration guidelines, sufficient space/room for preparation of medication, educational programs and continuous training for health care providers.
- Nursing managers apply of strategies in work such as clinical pharmacists, and computer/information technology.
- Nursing managers should establish anonymous reporting systems and counseling classes to create a secure environment to reduce nurses' fear and offer incentives to promote reporting.

Bibliography

1. Lisby M., *et al.* "Errors in the medication process: frequency, type, and potential clinical consequences". *International Journal for Quality in Health Care* (2005): 15-22.
2. WHO. "Patient safety: making health care safer" (2017).
3. WHO. "WHO launches global effort to halve medication-related errors in 5 years" (2017): 1.
4. National Coordinating Council for Medication Error Reporting and Prevention. "About medication errors" (2020).
5. Rahimi, F., *et al.* "Factors influencing medication errors according to nurses' decisions to do self-report". *International Journal of Basic and Clinical Pharmacology* (2015): 130-133.
6. Vilela B., *et al.* "Medication errors: management of the medication error indicator toward a more safety nursing practice". *Revista de Enfermagem UFPE* 10.1 (2016).
7. Zakharov S., *et al.* "Medication errors-an enduring problem for children and elderly patients". *Uppsala Journal of Medical Sciences* (2012): 309-317.
8. Ojerinde AC and Adejumo PO. "Factors associated with medication errors among health workers in University College Hospital, Nigeria". *Journal of Nursing and Health Science* (2014): 22-23.
9. Aboshaiqah AE. "Nurses' perception of medication administration errors". *American Journal of Nursing Research* (2014): 63-67.

10. Fathi A., *et al.* "Medication errors among nurses in teaching hospitals in the west of Iran: what we need to know about prevalence, types, and barriers to reporting". *Epidemiology and Health* (2017).
11. Miladinia M., *et al.* "Pediatric nurses' medication error: the self-reporting of frequency, types and causes". *International Journal of Pediatric* (2016): 1439-1444.
12. Johari H., *et al.* "Medication errors among nurses in government hospital". *Journal of Nurings and Health Sciences* (2013): 18-23.
13. Sears K and Goodman WM. "Risk factors for increased severity of paediatric medication administration errors". *Healthcare Policy* (2012): 109-126.
14. Abdullah D and Younus Sameen F. "Barriers that Preventing the Nursing Staff from Reporting Medication Errors in Kirkuk City Hospitals". *Kufa Journal for Nursing Sciences* (2017): 103-111.
15. Mrayyan., *et al.* "Rate, causes and reporting of medication errors in Jordan: nurses' perspectives". *Journal of Nursing Management* (2007): 659-670.
16. Lee E. "Reporting of medication administration errors by nurses in South Korean hospitals". *International Journal for Quality in Health Care* (2017): 728-734.
17. Yung HP, *et al.* "Nurses' attitudes and perceived barriers to the reporting of medication administration errors". *Journal of Nursing Management* (2016): 580-588.
18. Biftu BB., *et al.* "Medication administration error reporting and associated factors among nurses working at the University of Gondar referral hospital, Northwest Ethiopia, 2015". *Bio Med Central Nursing* (2016): 43.
19. Saleh AA and Barnard A. "Barriers Facing Nurses in Reporting Medication Administration Errors in Saudi Arabia". *American Journal of Nursing* (2019): 598-625.
20. Alarfj AA. "Barriers of Reporting Medication Errors at Arar Central Hospital, KSA". *Journal of Medical and Pharmaceutical Sciences* (2020): 36-56.
21. Mansouri SF, *et al.* "Barriers to nurses reporting errors and adverse events". *British Journal of Nursing* (2019): 690-695.
22. Mostafaei D., *et al.* "Medication errors of nurses and factors in refusal to report medication errors among nurses in a teaching medical center of Iran in 2012". *Iranian Red Crescent Medical Journal* (2014): e16600.
23. Mansi K., *et al.* "Biochemical Factors Relevant to Kidney Functions among Jordanian Children with Beta-thalassemia Major Treated with Deferoxamine". *International Journal of Medicine and Medical Sciences* (2013): 374-379.
24. ALBashtawy M., *et al.* "Self - medication among school students". *Journal of School Nursing* (2015): 110-116.
25. Alkhalwaldeh A., *et al.* "Long-term conditions associated with primary health care service use among older adults". *Primary Health Care* (2016): 31-35.
26. Sulaiman M., *et al.* "Exploring Safety Culture in Jordanian Hospitals: A Baseline Study". *Journal of Nursing Care Quality* (2017): E1-E7.
27. Alhaliqa F, *et al.* "Adherence to antipsychotic drug measured by an Arabic version of Morisky scale". *Issues in Mental Health Nursing* (2016): 975-982.

28. Qaddumi J., *et al.* "Prevalence of Hypertension and Pre-hypertension among Secondary School Students". *International Journal of Advanced Nursing Studies* (2016): 240-245.
29. ALBashtawy M., *et al.* "Assessment of headache among high school students in Jordan". *The Journal of School Nursing* (2017): 88-95.
30. Khatatbeh M., *et al.* "Prevalence of nicotine dependence among university students in Jordan: a cross-sectional study". *Epidemiology Biostatistics and Public Health* (2019).
31. Abdullah Al Kazaleh and Mohammed AL Bashtawy. "Therapeutic Communication Skills in Nursing Education and Practice". *EC Psychology and Psychiatry* 8.12 (2019): 01-04.

Volume 9 Issue 8 August 2020

©All rights reserved by Mohammad ALBashtawy, *et al.*