

Mechanical Power and the use of Mechanical Ventilation in Pediatric Acute Respiratory Distress Syndrome

José Colleti Junior^{1*} and Werther Brunow de Carvalho²

¹Pediatric ICU, Department of Pediatrics, Hospital Santa Catarina, São Paulo, Brazil ²Instituto da Criança, Department of Pediatrics, University of São Paulo, Brazil

*Corresponding Author: José Colleti Junior, Pediatric ICU, Department of Pediatrics, Hospital Santa Catarina, São Paulo, Brazil.

Received: August 08, 2017; Published: August 23, 2017

Children undergoing mechanical ventilation (MV) due to acute respiratory distress syndrome (ARDS) has been a challenge in the pediatric ICU, and although many advances have been done, there is still a high mortality (18% to 27%) [1]. Many guidelines and consensus have been published in recent years [1,2]; however, there is a lot of research to be done in this field to clarify the best practices, since the evidences in pediatric ARDS are still weak.

The guidelines focus on the protective ventilation, suggesting low tidal volumes (VT), set according to ideal body weight (IBW) of the patient to avoid barotrauma, volutrauma and ventilator-induced lung injury (VILI) [1,2]. However, all these recommendations are extrapolations from adult studies [3] and recent studies have shown that ARDS patients who are ventilated according to these guidelines may still be exposed to forces that can induce or aggravate lung injury [4,5].

Amato., *et al.* published a study in NEJM demonstrating that the driving pressure (ΔP) was associated with survival in adult patients with ARDS [6]. He concluded that "decreases in ΔP owing to changes in ventilator settings were strongly associated with increased survival". That was another step in the knowledge on how to ventilate patients with ARDS, and although there are no similar studies in children, we are again extrapolating data from this study, and using them to ventilate pediatric patients with ARDS. However, we should remember that the pediatric patient is not a "mini-adult" and has its particularities. Many treatments that are well-established for adults are useless or harmful to children. Therefore, we need clinical trials to validate these data in children.

More recently, the concept of mechanical power has been raised as a more accurate formula to predict VILI [7], since the current strategy to minimize VILI has been to avoid high VT or ΔP ; however, the role of "power" (work applied to the lung per unit time) is uncertain. Hence, other variables are taken into account, besides the VT and the ΔP . Then the mechanical power equation may help estimate the contribution of the different ventilator-related causes of lung injury and of their variations.

Although these data are from adult patients, the general idea is quite intuitive and should be translated to the pediatric patients, while awaiting the clinical trials.

Competing Interests

The authors declare that they have no competing interests.

Grants

The authors declare that no grants were involved in supporting this work.

Bibliography

1. Khemani RG., *et al.* "Pediatric acute respiratory distress syndrome: definition, incidence, and epidemiology: proceedings from the Pediatric Acute Lung Injury Consensus Conference". *Pediatric Critical Care Medicine* 16 (2015): S23-S40.

Citation: José Colleti Junior and Werther Brunow de Carvalho. "Mechanical Power and the use of Mechanical Ventilation in Pediatric Acute Respiratory Distress Syndrome". *EC Pulmonology and Respiratory Medicine* 4.6 (2017): 176-177.

Mechanical Power and the use of Mechanical Ventilation in Pediatric Acute Respiratory Distress Syndrome

- 2. Pediatric Acute Lung Injury Consensus Conference Group. "Pediatric acute respiratory distress syndrome: consensus recommendations from the Pediatric Acute Lung Injury Consensus Conference". *Pediatric Critical Care Medicine* 16.5 (2015): 428-439.
- 3. Amato MB., *et al.* "Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome". *New England Journal of Medicine* 338.6 (1998): 347-354.
- 4. Terragni PP., *et al.* "Tidal hyperinflation during low tidal volume ventilation in acute respiratory distress syndrome". *American Journal of Respiratory and Critical Care Medicine* 175.2 (2007): 160-166.
- 5. Grasso S, *et al.* "ARDSnet ventilatory protocol and alveolar hyperinflation: role of positive end-expiratory pressure". *American Journal of Respiratory and Critical Care Medicine* 176.8 (2007): 761-767.
- Amato MB., *et al.* "Driving pressure and survival in the acute respiratory distress syndrome". *New England Journal of Medicine* 372.8 (2015): 747-755.
- 7. Cressoni M., et al. "Mechanical Power and Development of Ventilator-induced Lung Injury". Anesthesiology 124.5 (2016): 1100-1108.

Volume 4 Issue 6 August 2017 ©All rights reserved by José Colleti Junior and Werther Brunow de Carvalho.