

## COVID-19 Influencing Thyroid Dysfunction

**Porntep Siriwanarangsun<sup>1</sup>, Attapon Cheepsattayakorn<sup>1,2\*</sup> and Ruangrong Cheepsattayakorn<sup>3</sup>**

<sup>1</sup>Faculty of Medicine, Western University, Pathumtani Province, Thailand

<sup>2</sup>10<sup>th</sup> Zonal Tuberculosis and Chest Disease Center, Chiang Mai, Thailand

<sup>3</sup>Department of Pathology, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand

**\*Corresponding Author:** Attapon Cheepsattayakorn, 10<sup>th</sup> Zonal Tuberculosis and Chest Disease Center, Chiang Mai, Thailand.

**Received:** December 15, 2020; **Published:** August 14, 2024

Recently, a first case-report of thyroid dysfunction following SARS-CoV-2 (COVID-19) infection, namely subacute thyroiditis was reported [1]. This female patient had a painful, enlarged thyroid gland [2]. The exact mechanisms that SARS-CoV-2 (COVID-19) causes thyroid dysfunction are not known. Nevertheless, mechanisms that are demonstrated through SARS virus are potential central mechanism [3-5], direct viral replication [6-9], interaction with thyroid-ACE2 receptor [10-12], and inflammatory response, apoptosis, and local damage [13-16]. SARS-CoV-2 (COVID-19) and thyroid dysfunction impact each other by: 1) Graves's ophthalmopathy with actively undergoing immunosuppressive therapy are likely to increase risk of severe coronavirus infection development [17]; 2) Patients with poorly controlled thyroid dysfunction, particularly those with thyrotoxicosis, may be at risk of thyroid storm [18]; and 3) Systemic disease, including COVID-19 are related to low-T3 syndrome or non-thyroidal illness [19]. As pregnant women with hyper- or hypothyroidism are at increased risk of development of more severe COVID-19 disease, they are particularly suggested to social distancing adherence [20]. In the first trimester of pregnancy, the preferred treatment is the lowest possible dose of propylthiouracil (PTU) [21].

In conclusion, the following endocrine service is suggested in the COVID-19 crisis: satellite blood-testing services, remote monitoring services, face-to-face appointments, and telephone and video consultation.

### Bibliography

1. Brancatella A., et al. "Subacute thyroiditis after SARS-CoV-2 infection". *The Journal of Clinical Endocrinology and Metabolism* 105 (2020): 2367-2370.
2. Guimarães VC. "Subacute and Riedel's thyroiditis". In: *Endocrinology: Adult and Pediatric*, 7<sup>th</sup> edition. Jameson JL, De Groot LJ, eds. Elsevier: Saunders, PA, USA (2016): 1541-1556.
3. Wang W, et al. "Evaluation and observation of serum thyroid hormone and parathyroid hormone in patients with severe acute respiratory syndrome". *The Journal of the Chinese Antituberculosis Association* 25 (2003): 232-234.
4. Leow MK, et al. "Hypocortisolism in survivors of severe acute respiratory syndrome (SARS)". *Clinical Endocrinology* 63.2 (2005): 197-202.
5. Chrousos GP and Kaltsas G. "Post-SARS sickness syndrome manifestations and endocrinopathy: how, why, and so what?" *Clinical Endocrinology* 63.4 (2005): 363-365.
6. Desailoud R and Hober D. "Viruses and thyroiditis: an update". *Virology Journal* 6 (2009): 5.
7. Chang L, et al. "Coronavirus disease 2019: coronaviruses and blood safety". *Transfusion Medicine Reviews* 34.2 (2020): 75-80.

8. Ding Y, *et al.* "Organ distribution of severe acute respiratory syndrome (SARS) associated coronavirus (SARS-CoV) in SARS patients: implications for pathogenesis and virus transmission pathways". *The Journal of Pathology* 203.2 (2004): 622-630.
9. Gu J, *et al.* "Multiple organ infection and the pathogenesis of SARS". *Journal of Experimental Medicine* 202.3 (2005): 415-424.
10. Liu F, *et al.* "ACE2 expression in pancreas may cause pancreatic damage after SARS-CoV-2 infection". *Clinical Gastroenterology and Hepatology* (2020).
11. Li MY, *et al.* "Expression of the SARS-CoV-2 cell receptor gene ACE2 in a wide variety of human tissues". *Infectious Diseases of Poverty* 9.1 (2020): 45.
12. Kuba K, *et al.* "A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury". *Nature Medicine* 11.8 (2005): 875-879.
13. Perlman S and Dandekar AA. "Immunopathogenesis of coronavirus infections: implications for SARS". *Nature Reviews Immunology* 5.12 (2005): 917-927.
14. Law PT, *et al.* "The 3a protein of severe acute respiratory syndrome-associated coronavirus induces apoptosis in Vero E6 cells". *Journal of General Virology* 86.7 (2005): 1921-1930.
15. Tan YJ, *et al.* "Overexpression of 7a, a protein specifically encoded by the severe acute respiratory syndrome coronavirus, induces apoptosis via a caspase-dependent pathway". *Virology Journal* 78.24 (2004): 14043-14047.
16. Yuan X, *et al.* "G0/G1 arrest and apoptosis induced by SARS-CoV 3b protein in transfected cells". *Virology Journal* 2 (2005): 66.
17. Antonelli A, *et al.* "Graves' disease: epidemiology, genetic and environmental risk factors and viruses". *Best Practice and Research: Clinical Endocrinology and Metabolism* 34 (2020): 101387.
18. De Leo S, *et al.* "Hyperthyroidism". *Lancet* 388 (2016): 906-918.
19. Fliers E, *et al.* "Thyroid function in critically ill patients". *Lancet: Diabetes and Endocrinology* 3 (2015): 816-825.
20. <http://www.rcog.org.uk/globalassets/documents/guidelines/2020-04-17-coronavirus-covid-19-infection-in-pregnancy.pdf>
21. Alexander EK, *et al.* "2017 Guidelines of the American Thyroid Association for the diagnosis and management of thyroid disease during pregnancy and postpartum". *Thyroid* 27 (2017): 315-389.

**Volume 13 Issue 8 August 2024**

**© All rights reserved by Attapon Cheepsattayakorn., *et al.***